



Effect of IBA Concentration (Indol Butaric Acid) and Growing Condition on Performance of Cuttings in Mogra (*Jasminum sambac*)

R. J. Makwana^{a++*}, V. D. Rathva^{a++} and B. H. Panchal^{b#}

^a Sheth D. M. Polytechnic in Horticulture Model Farm, AAU, Vadodara 390003, India.

^b Krishi Vigyan Kendra, AAU, Arnej, Gujarat, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i102998>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/125672>

Original Research Article

Received: 15/08/2024

Accepted: 19/10/2024

Published: 28/10/2024

ABSTRACT

An experiment was laid out to study influence of different levels of IBA and growing condition on cuttings of Mogra (*Jasminum sambac*) at Sheth D. M. polytechnic in Horticulture College, Anand Agricultural University, Model farm, Vadodara during 2022 to 2024. The study was conducted using completely randomized design with factorial concept and three replications for each treatment. Total 10 treatment combination of two factors viz., growing condition (open and NVPH) and IBA concentration (500, 1000, 1500 and 2000 ppm) were compared with control (deepen in distilled water). IBA significantly improved the performance of cuttings under naturally ventilated poly-house

⁺⁺Assistant Professor;

[#]Senior Scientist and Head;

^{*}Corresponding author: E-mail: drshimak@aaui.in;

Cite as: Makwana, R. J., V. D. Rathva, and B. H. Panchal. 2024. "Effect of IBA Concentration (Indol Butaric Acid) and Growing Condition on Performance of Cuttings in Mogra (*Jasminum Sambac*)". *Journal of Experimental Agriculture International* 46 (10):744-50. <https://doi.org/10.9734/jeai/2024/v46i102998>.

condition. 1500 ppm IBA concentration recorded significantly lower days to sprout (13.83 days), higher shoot length at 30, 45 and 60 DAP (2.80 cm, 7.64 cm and 11.63 cm), root length (4.18 cm), number of leaves at 30, 45 and 60 DAP (2.87, 6.03 and 8.47), fresh weight (11.63g), dry weight (4.18g) and survival percentage (66.11%). Where as in growing condition, naturally ventilated poly-house recorded significantly lower days to sprout (13.27 days), higher shoot length at 30, 45 and 60 DAP (2.40cm, 7.26 cm and 11.80 cm), root length (3.89 cm), number of leaves at 30, 45 and 60 DAP (2.72 , 4.80 and 7.85), fresh weight (11.80 g), dry weight (3.89 g) and survival percentage (64.56%). Semi hardwood cuttings of *Jasminum sambac* treated with 1500 ppm IBA concentration treatment for 5 min and planted in polybags under Naturally ventilated poly-house showed promising growth with higher sprouting and survival percentage as compared to other treatments. It was recommended to propagate semi hardwood cuttings of *Jasminum sambac* with 1500 ppm IBA concentration for 5 min and plant them in naturally ventilated polyhouse to get maximum sprouting percentage and quality planting material.

Keywords: *Jasminum sambac*; cuttings; IBA; Naturally ventilated poly-house; plant growth regulators.

1. INTRODUCTION

In India, jasmines are cultivated throughout the country but the commercial cultivation is confined to Coimbatore, Madurai, and Dindigul (Tamil Nadu); Bangalore, Bellary, Mysore and Kolar (Karnataka); Knnauj, Jaunpur and Gazipur (Uttar Pradesh); Udaipur, Jaipur, Ajmer and Kota (Rajasthan); Ranaghat, Kolaghat, Pancskura (West Bengal); parts of Andhra Pradesh, Maharashtra and Gujarat. Gujarat has established itself as a major companion in the floriculture sector with about 22 thousand ha area under flower crops and approximately 201 Thousand MT of production [1]. In *Jasminum sambac* seeds are not formed therefore the vegetative propagation is the only reproductive method. Normally vegetative propagation is achieved through ground layering but it is not convenient for transportation purpose of germplasm [2]. Pruning is the most important practice in jasmine that can enhance the health, vigour and aesthetic of Mogra plant. The left over material of pruned benches may become easy source of vegetative propagation to produce new plants and also generating extra income for farmers. Rushabh *et al.* (2024) studied influence of time of pruning on flower yield and observed that pruning during December month gives higher yield in Jasmine (*Jasminum sambac*). Cutting is the most popular way of multiplication of ornamental shrubs [3], but rooting success rate through conventional method of hardwood cuttings is very low. Renuka and Sekhar [4] studied the influence of different concentrations of IBA on carnation cv. Dona and recorded highest percentage of establishment of rooted cuttings at 200 ppm IBA. Similar finding have Similar findings were observed by Hirapara [5] in Jasmine. Hence an experiment is been laid out

to optimise best IBA concentration and growing conditions for cutting in *Jasminum sambac* cv. Local.

2. MATERIALS AND METHODS

The present investigation was carried out during 2022 and 2023 at Sheth D. M. Polytechnic in Horticulture college research farm, Anand Agricultural University, Model farm, Vadodara to study the effect of levels of IBA (Indol butaric acid) and growing condition on performance of cuttings in Mogra (*Jasminum sambac*) with ten treatment combination of two factor *viz.*, growing condition (Open and Naturally ventilated poly-house) and level of IBA (500, 1000, 1500 and 2000 ppm) in completely randomized design with factorial concept and three replications for each treatment. Semi hardwood cuttings of *Janminum sambac* were collected on second week of December from 3 year old plants having 25-30 cm long cuttings with 5 to 8 nodes. 60 cutting for each treatment was selected and basal end of 2-3 cm portion of the cuttings was dipped in growth regulator formulation according to the treatment for 5 minutes and divided in 20 cuttings per replication and immediately planted in polybags containing prescribed media to a depth of 6-8 cm and placed under NVPH condition and another set of same treatment was placed in open condition. The cuttings under control were dipped in distilled water instead of plant growth regulators. Planted cuttings were treated with 1 per cent Bavistin to prevent the occurrence of fungal diseases. IBA was diluted in 1N NaOH and growing media was prepared with mixing the well sieved sand, soil and vermicompost (1:2:1) in polybags. Various observations like days to sprouting, plant height at 30, 45 and 60 days after planting (DAP), number of leaves at 30, 45

and 60 days after planting root length, fresh weight, dry weight and survival percentage at were taken during the experiment period. The experiment was conducted for two years and pooled data was taken to statistical analysis by the method proposed by Panse and Sukhatme [6].

3. RESULTS AND DISCUSSION

From the data depicted in Tables 1 and 2 revealed that Level of IBA and growing condition influenced the seedling parameters viz., days to sprouting, plant height at 30, 45 and 60 DAP, number of leaves at 30, 45 and 60 DAP, length of roots, fresh weight and dry weight and survival percentage while interaction was found significant for length of roots, fresh weight, dry weight, survival percentage, shoot length at 30 and 60 DAP and number of leaves at 30 DAP whereas, interaction effect was found non-significant for days to sprouting.

3.1 Effect of IBA

3.1.1 Days to Sprouting

In case of level of IBA, cutting treated with 2000 ppm IBA treatment T5, recorded significantly lower days to sprouting (13.75 days) which was at par with 1500 ppm IBA treatment T4 (13.83 days) whereas, maximum days to sprouting was recorded in control (17.75 days). This is may be due to the perfect amount of plant growth regulator and its concentration, which enhance the process of cell division, cell elongation and early differentiation of callus tissue toward the root formation ultimately resulted early growth of cuttings. In vegetative propagation, early differentiation and growth of shoot largely relies on food reserves available in the cuttings [7]. This is followed by early root formation, which ultimately absorbs more nutrients and water there by encourage sprouting of shoot. Similar trend of finding was also confirmed by Netam *et al.* [8], Santhosha *et al.* [9], Nagaraja *et al.* [10] and Pooja and Sadatulla [11] in Jasmine and Chovatia *et al.* [12] in Bougainvillea, Renuka and Sekhar [4] in Carnation.

3.1.2 Length of shoot

In cuttings length of the vegetative shoot is one of the main character which represents the vegetative growth of cuttings. In experiment, length of shoot was recorded at 30, 45 and 60 days after planting of cuttings. The data clearly

showed that cutting treated with 1500 ppm IBA concentration treatment T4 recorded significantly higher shoot length at 30, 45 and 60 DAP (2.80 cm, 7.64 cm and 11.63 cm respectively) which was at par with 2000 ppm concentration of IBA treatment T5 at 30 and 60 days after planting. The significant difference in length of shoot might be due to early sprouting of vegetative buds in IBA treated cuttings. Auxin enhanced cell division and cell enlargement, promotion of protein synthesis which might have resulted in enhanced vegetative growth [13]. Similar findings were observed by Nagaraja *et al.* [10], Pooja and Sadatulla [11], Netam *et al.* [8] and Sharma and Brar [2] in Jasmine.

3.1.3 Length of Roots

Data presented in Table 1 showed that cuttings treated with 1500 ppm IBA treatment T4, recorded maximum length of roots per cuttings (4.18 cm) which was at par with cuttings treated with 2000ppm IBA treatment T5 (4.15 cm). Similar findings were observed by Santhosha *et al.* [9], Netam *et al.* [8], Zaghoul *et al.* [14] and Patil *et al.* [15] in Jasmine and Renuka and Sekhar [4] in Carnation.

3.1.4 Number of leaves

The number of leaves were recorded at time interval of 30, 45 and 60 DAP of cuttings in this experiment. Data presented in Table 2 recorded cuttings treated with 1500 ppm IBA treatment T4 recorded highest number of leaves at 30, 45 and 60 DAP (2.87, 6.03 and 8.47 respectively), which was at par with cuttings treated with 2000 ppm IBA at 30 and 60 DAP. Increase in number of leaf in shoot might be due to vigorous growth and early initiation of root induced by the application of growth regulator which helps to absorbs more nutrients and thereby producing more leaves as reported by Stancato *et al.* [16]. Similar findings were observed by Netam *et al.* [8], Pooja and Sadatulla [11] and Hirapara [5] in Jasmine.

3.1.5 Fresh Weight and Dry Weight

From the data depicted in Table 2, it was evident that cutting that was deepen in 1500 ppm IBA solution treatment T4 recorded significantly higher fresh weight (11.63 g/plant) and dry weight (4.18 g/plant) which was at par with cutting treated with 2000 ppm IBA solution treatment T5. It might be due early and fast cell division and cell enlargement process with early and easy initiation of roots due to presence of

Table 1. Effect of levels of IBA and growing condition on days to sprouting (days), shoot length at 30, 45 and 60 DAP and length of roots at 60 DAP in Jasmine (*Jasminum sambac*)

Treatments	Days to sprouting	Shoot length at (cm)			Length of roots (cm)
		30 DAP	45 DAP	60 DAP	
Factor A: Growing conditions					
G ₁ : Open Conditions	17.87	2.03	5.95	9.13	3.01
G ₂ : NVPH	13.27	2.40	7.26	11.80	3.89
C.D. (p=0.05)	0.46	0.09	0.25	0.34	0.17
Factor B: IBA Concentrations					
T ₁ : Distill water	17.75	1.52	5.59	9.01	2.65
T ₂ : 500 ppm	16.58	1.98	6.36	9.81	2.82
T ₃ : 1000 ppm	15.92	2.01	6.68	10.31	3.46
T ₄ : 1500 ppm	13.83	2.80	7.64	11.63	4.18
T ₅ : 2000 ppm	13.75	2.76	6.74	11.55	4.15
C.D. (p=0.05)	0.72	0.14	0.39	0.54	0.26
Interaction effect (G x T)	NS	Sig	NS	Sig	Sig

Table 2. Effect of levels of IBA and growing condition on number of leaves at 30, 45 and 60 DAP, fresh weight, dry weight and survival percentage at 60 DAP in Jasmine (*Jasminum sambac*)

Treatments	Number of leaves			Fresh weight (g/plant)	Dry weight (g/plant)	Survival percentage (%)
	30 DAP	45 DAP	60 DAP			
Factor A: Growing conditions						
G ₁ : Open Conditions	2.27	4.32	6.14	9.13	3.01	46.00
G ₂ : NVPH	2.72	4.80	7.85	11.80	3.89	64.56
C.D. (p=0.05)	0.13	0.09	0.30	0.34	0.17	2.49
Factor B: IBA Concentrations						
T ₁ : Distill water	2.20	3.28	5.45	9.01	2.65	39.72
T ₂ : 500 ppm	2.30	3.55	6.00	9.81	2.82	51.67
T ₃ : 1000 ppm	2.42	4.45	7.02	10.31	3.46	53.61
T ₄ : 1500 ppm	2.87	6.03	8.47	11.63	4.18	66.11
T ₅ : 2000 ppm	2.70	5.48	8.03	11.55	4.15	65.28
C.D. (p=0.05)	0.20	0.30	0.48	0.54	0.26	3.94
	Sig	NS	NS	Sig	Sig	Sig

auxin. Fresh matter accumulation of plant depends upon the vegetative growth parameter viz., plant height, spread or average number of branches and it is directly influenced by auxin. Thus it enhanced the fresh matter accumulation, resulted from photosynthesis or the hydrolysis of starch resulting in increase in concentration of sugar [17]. The results are in line with the findings observed by Girisha *et al.* [18] in Daisy and Netam *et al.* [8] and Chaitnaya (2013) in Jasmine.

3.1.6 Survival Percentage

Survival percentage is ultimate observation to check the final outcome of the cutting during experimental period. Cutting treated with 1500 ppm IBA solution recorded highest survival

percentage (66.11 %) which was at par with treatment T5. Increase in survivability of cutting may be due to initiation of more roots and number of leaves with IBA treatments. The finding agrees with the finding of Chowdhuri *et al.* [19], Pooja and Sadatulla [11] and Shanthosha *et al.* [9] in jasmine.

3.1.7 Effect of Growing Condition

Naturally ventilated poly-house provide favorable conditions for plants as well as newly developed cutting for better growth. Application of plant growth regulators under favorable growing conditions significantly improve the performance of the cuttings in rose [20]. Present investigation was conducted to compare open field condition (G1) with the naturally ventilated poly-house

(G2). Growing condition showed significant influence on all the parameters over open field condition (control). Cuttings placed under naturally ventilated poly-house recorded significantly lower days to sprout (13.27 days), higher shoot length at 30, 45 and 60 DAP (2.40cm, 7.26 cm and 11.80 cm), root length (3.89 cm), number of leaves at 30, 45 and 60 DAP (2.72 , 4.80 and 7.85), fresh weight (11.80 g), dry weight (3.89 g) and survival percentage (64.56%). It was evident cuttings performed well

under naturally ventilated poly-house that might be due to better favorable condition to grow. Poly-hose decreases 66% global solar intensity which becomes advantage to cutting for better growth. Solar radiation is a determining factor of the climatic variable inside the protected structures. Micro climate including temperature, relative humidity, moisture etc plays important role for growth of cuttings. These results are in agreement with the findings of Ahmed *et al.*, [21] and Chakir *et al.*, [22-24].



Fig. 1. Effect of IBA on semi hard wood cuttings of Jasmine (*Jasminum sambac*) kept under open condition at 60 days after planting



Fig. 2. Effect of IBA on semi hard wood cuttings of Jasmine (*Jasminum sambac*) kept under NVPH condition at 60 days after planting

4. CONCLUSION

It was concluded from the present investigation that *Jasminum sambac* var. Local cutting treated with 1500 ppm IBA concentrated solution for 5 minutes provided better growth with early initiation and higher survival percentage under naturally ventilated poly-house condition.

5. FUTURE SCOPE

Jasmine cutting are difficult to root and there are several species of jasmine which have different growth habit and response to propagated by cuttings. More research in terms of response under various month and season effect can be taken to understand the response of cuttings in jasmine crop to different kind of plant growth regulators.

6. DECLARATION

We believe that our findings could be of interest to the scientist, farmers, nurseryman and reader of the journal. We certainly certify about this article is an original research work and the same is not submitted or published earlier elsewhere. We declare that there are no conflicts of interests associated to this manuscript. Additionally, this manuscript meets all applicable standards of an agricultural research with regard to the ethics of experimentation and research work; also there is no duplicate publication, fraud, plagiarism, or concerns about animal or human experimentation during the research work.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Department of horticulture annual report. 2023-24
Available:<https://doh.gujarat.gov.in/Home/HorticultureCultivation..>

2. Sharma SP, Brar RS. High frequency multiplication of jasmine (*Jasminum sambac* L.) Aiton using plant growth hormone solutions on stem cutting. International Journal of Applied Life Sciences and Engineering. 2014; 1(1):70-73.
3. Bose TK, Mukherjee TP, Roy T. Standardization of propagation from cutting under mist. *Punjab Hort. J.*, 1975;15(3-4):139-143.
4. Renuka K, Sekhar RC. Studies on effect of plant growth regulators on rooting of carnation (*Dianthus caryophyllus* L.) cuttings of cv. Dona under poly house conditions. *Plant Archives*. 2014;14(2): 1135-1137.
5. Hirapara DV. Effect of IBA and NAA on Vegetative Propagation of *Jasminum arborescens* L. Cv.'Paras' Through Semi-Hard Wood Cutting (Doctoral dissertation, Horticulture Department, NM College of Agriculture, Navsari Agriculture University); 2005.
6. Panse VG, Sukhatme PV. Statistical methods for agricultural workers. Publication and Information Division, ICAR, New Delhi; 1967.
7. Nanda KK. Physiology of adventitious root formation. *Ind. Jour. Plant Physiol.* 1975;18: 80–89.
8. Netam N, Shukla N, Sharma G, Sahu JK. Effect of Different IBA concentration on survivability and rooting of Jasmine (*Jasminum sambac* (L.) Aiton) stem cuttings. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(1S):614-617.
9. Santhosha HM, Patil R, Manju MJ. Study of floral biology and standardization of soft wood cutting propagation technique in Bhatkal jasmine (*Jasminum sambac* var Bhatkal Jasmine). *The Pharma Innovation Journal*. 2021;10(11): 1846-1850.
10. Nagaraja GS, Rai Muthappa BC, Guruprasad TR. Effect of Intermittent mist and growth regulators on propagation of *Jasminum grandiflour* by different types of cutting. *Haryana J. Wort. Sci*. 1991;20(3-4):183-188.
11. Pooja HS, Sadatulla F. Effect of Indole-3-Butyric Acid on Growth of V-1 Mulberry Cuttings with Varying Number of Buds in Bangalore. *International Journal of Plant & Soil Science*. 2022;34(24):9-13.
12. Chovatia VP, Poshia VK, Shukla PT. Root initiation studies in bougainvillea (*Bougainvillea peruviana* L.) var. Mary

- Palmer. G.A.U. Res. J. 1995;20(2):167-169
13. Evans ML. Rapid stimulation of plant cell elongation by hormonal and non-hormonal factors. *Bioscience*. 1973;23:7-8.
 14. Zaghoul MA, El R, Hassan SM, Elgindy WM. Effect of NAA and Kinetin on rooting of Jasmine sambac. *Annals of Agric. Sci*. 1990;35(2):939-447.
 15. Patil B, Kanamadi VC. Effect of growth regulators on the rooting of hardwood cuttings in jasmine. *Advances in Forestry Research in India*. 1998;19:225-229.
 16. Stancato GC, Aguiar FFA, Kanashiro S, Tavares AR, Catharino ELM, Silveira RBDA. *Rhipsalis grandiflora* Haw. (Cactaceae) propagation by stem cuttings. *Scientia Agricola*. 2003;60:651-656.
 17. Thimann K. The natural plant hormones. *Plant physiology, a Treatise*. 1972;3:365.
 18. Girisha R, Shirol AM, Reddy BS, Patil VK, Krishnamurthy GH. Growth, quality and yield characteristics of daisy (*Aster amellus* L.) cultivar Dwarf Pink as influenced by different plant growth regulators. *Karnataka J. Agric. Sci*. 2012;25(1):163-165.
 19. Chowdhuri TK, Sadhukhan R, Mondal T, Das S. Effect of different growth regulators on propagation of Cape jasmine (*Tabernaemontana coronaria* var. Dwarf) in subtropical zone of West Bengal. *The Asian journal of horticulture* 2017;12(2):206-210.
 20. Kentelky E, Székely-varga Z, Balla G. Propagation of rose varieties by cuttings under the effect of different rooting hormones. *Acta Universitatis Sapientiae Agriculture And Environment*, 2023;15:98–109
 21. Ahmed HA, Tong Y, Yang Q, Al-Faraj AA, Abdel-Ghany AM. Spatial distribution of air temperature and relative humidity in the greenhouse as affected by external shading in arid climates. *J Integr Agric*. 2019;18(12):2869-82.
 22. Chakir S, Bekraoui A, Majdoubi H, Mouqallid M. Microclimate and Efficiency Analysis of an Olive Rooting Cuttings Tunnel Greenhouse Type under Mediterranean Wintertime Climatic Conditions. *The Open Agriculture Journal*. 2022;16(1):1-13
 23. Poorna CR, Chandrasekhararao C, Suchitra V, Salomi Suneetha DR. Effect of IBA concentration and size of cutting on rooting of *Jasminum sambac*. *Green Farming*. 2015;6(4):888-892.
 24. Makwana RJ, Rathva VD, Mahida SV. Effect of different level and time of pruning on growth and flowering in Mogra (*Jasminum sambac*) var. Local. *International Journal of Research in Agronomy*. 2024;7(1):424-427

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/125672>