



# Proximate, Phytochemical and Mineral Composition of Methanol Extract of *Persea americana* Seed

Omorovbiye Omorevbarhia<sup>a\*</sup>, Sunday Chidozie Umerie<sup>b</sup>,  
Ejike Louis Aniefuna<sup>c</sup> and Oguazu Chinenye Enoch<sup>b</sup>

<sup>a</sup> Department of Biochemistry, Faculty of Natural Sciences, Paul University, Awka, Anambra State, Nigeria.

<sup>b</sup> Department of Applied Biochemistry, Faculty of Biosciences, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria.

<sup>c</sup> Department of Mineral Resources and Energy, Ministry of Petroleum and Mineral Resources, Awka, Anambra State, Nigeria.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJBCRR/2023/v32i2797

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97536>

Original Research Article

Received: 16/01/2023

Accepted: 18/03/2023

Published: 29/03/2023

## ABSTRACT

This study investigated the proximate composition and evaluated the phytochemicals present in the methanol extract of *Persea americana* seed. The seed samples were collected, dried, ground into powder, and extracted in methanol by cold maceration. Proximate analysis was carried out using the standard methods of association of official analytical chemists (AOAC) while quantitative phytochemical analysis was done using standard methods. Mineral composition assays were carried out using the standard methods of the American public health association (APHA). Proximate analysis revealed that the seeds of *P. americana* contain more carbohydrate (32.05±1.21%), fat (25.73±2.01%), and protein (21.00±0.88%) with fewer amounts of moisture

\*Corresponding author: E-mail: omorevbarhiao14008f@gmail.com;

(8.98±0.10%), and ash (3.28±0.16%). Quantitative phytochemical analysis revealed the presence of phenols (3.51±0.42 mg/100 g), saponins (0.05±0.38 mg/100g), alkaloid (0.70±0.37 mg/100g), beta-carotene (32.24±0.55 mg/100 g), lycopene (6.84±0.52 mg/100 g), flavonoids (34.14±3.13 mg/100 g), phytate (0.00±0.03 mg/100 g), cardiac glycosides (0.37±0.01%), total phytosterol (21.25±0.12 mg/100 g), stigmasterol (13.16±0.15mg/100g) and campesterol (6.16±0.12 mg/100 g). Mineral contents of the sample revealed that potassium, calcium, and magnesium have the highest concentrations while manganese and iron have the lowest. The chemical compositions of the investigated samples might be responsible for their medicinal values in phytomedicine. This study shows that *Persea americana* seeds are adequate in lowering cholesterol levels probably due to the presence of phytosterols.

**Keywords:** *Persea americana* seeds; phytochemical composition; proximate; minerals analysis.

## 1. INTRODUCTION

*Persea americana* Mill. (Lauraceae), a plant from Central America (Mexico, Guatemala, Antilles), has shown easy adaptation to other tropical regions including Africa [1]. "The fruit is popularly known as avocado pear and has an olive-green peel and thick pale yellow pulp that is rich in vegetable oils and appreciated for its sensory attributes" [1]. Avocado pear pulp is consumed similarly to the African pear fruit pulp (*Dacryodes edulis*). However, despite the nutritional potential of African pear fruit seed [2], it is still discarded after the edible pulp is eaten [3]. "The avocado seed represents 13–18% of the fruit and is a byproduct generally not utilized" [4]. Normally, the seed is discarded during the processing of the pulp like the African pear fruit seed. Discarding the seed may cause severe ecological problems [4]; like increased numbers of insects and rodents and economic losses as a result of the high cost of transporting these byproducts to disposal areas [5].

The seed of *P. americana* has been reported to lower blood pressure [6] with a reduction in the total cholesterol, LDLC (low-density lipoproteins cholesterol), and triacylglycerol in the plasma, kidney, liver, and heart of normotensive and hypertensive rat models at high doses of the seed extract [7]. "In hypercholesterolemia, a condition of high blood cholesterol, LDLC does not aid the transportation of cholesterol out of body tissues but deposits cholesterol in the lumen of arterial walls" [8]. "Phytosterols have been found effective in treating high cholesterol (hypercholesterolemia) as the plant sterols compete for absorption sites with cholesterol; they thus reduce the amount of cholesterol absorbed" [9]. "The cholesterol-lowering effect of phytosterols is also believed to be caused by an inhibition of cholesterol absorption resulting from the higher solubility of phytosterols than of

cholesterol in bile salt micelles. The most common phytosterols are campesterol and stigmasterol which occurs in higher plants both *in vivo* and *in vitro* tissue culture" [9].

Meanwhile, Joao et al., [1] and Arukwe et al., [10] have shown that avocado seeds contain bioactive compounds such as flavonoids, phenols, alkaloids, saponins, and phytosterols that have tremendous health benefits. These compounds have antioxidant properties that help in preventing and treating putative health diseases such as cancer, atherosclerosis, diabetes, hypertension, Alzheimer's disease, and ulcer [11].

## 2. MATERIALS AND METHODS

### 2.1 Sample Collection and Identification

Avocado pear samples were purchased from Eke Awka Market in Awka, Anambra State. The pear samples were identified by a taxonomist in the Department of Botany, Nnamdi Azikiwe University Awka, Anambra State.

### 2.2 Preparation of Sample Extract

The seeds were separated from the fruit, washed, chopped in bits, and then dried at room temperature for ten days and then ground into a fine powder using a manual grinder. The powdered sample was stored in an air-tight container until further analysis. The sample extraction was done by cold maceration as described by Kumar et al., [12]. The powdered sample (10g) was dissolved in 100% methanol (100 ml) for 24 hours. The mixture was filtered through Whatman paper No. 4 and the filtrate was concentrated over a water bath at 40°C. The concentrated extract was weighed and redissolved in methanol at a concentration of 100mg/ml and stored at 4°C for further analysis.

## 2.3 Proximate Analysis

Moisture content, ash content, crude fiber, protein, and fat were determined using standard methods of AOAC, [13] while total carbohydrate content was determined by a difference of 100 as described by Merrill and Watt [14].

## 2.4 Qualitative Analysis of Phytochemical Constituents

Qualitative analysis of the phytochemicals (alkaloids, flavonoids, tannins, saponins, phenols, and terpenoids) of *Persea americana* seeds was carried out using the methods of Trease and Evans, [15] and Harborne [16].

## 2.5 Quantitative Determination of Phytochemical Constituents

The phytate and oxalate contents were determined by titration using the methods of Young and Greaves [17] and Osagie, [18] respectively. Terpene was determined according to the method of Narayan et al., [19]. The saponins and alkaloids content was weighed and

calculated in percentage [16]. The tannin content was calculated and expressed in percentage (Follins Dennis titration method) as described by Pearson [20]. Beta carotene and lycopene, total phenol, and flavonoid contents of the sample were determined using the method of Barros et al., [21]. The phytosterol content of the sample was determined according to the method of Larissa et al., [22] while mineral content was done using Varian AA240 Atomic Absorption Spectrophotometer according to the method of APHA [23].

## 3. RESULTS

The seeds of *Persea americana* were investigated to evaluate the proximate, phytochemical, and mineral compositions, the results obtained are as follows:

### 3.1 Results for Proximate Analysis

This result shows that *Persea americana* seeds are rich in carbohydrates and lipid but the ash content is relatively low.




Fig. 1. Bar chart of proximate composition of *Persea americana* seed

### 3.2 Results for Phytochemical Composition

This result shows that *Persea americana* seeds are highly packed with tannins, terpenoids but low in phytate, alkaloids and saponins.

### 3.3 Results for Mineral Content

This result shows that *Persea americana* seeds are relatively high in potassium, phosphorus, and magnesium but low in manganese, iron, and copper.

## 4. DISCUSSION

The seeds of *Persea americana* were investigated to evaluate their proximate, phytochemical, and mineral compositions.

The proximate composition of *Persea americana* seeds as shown in Fig. 1 revealed that the moisture content was relatively low and could imply long shelf life [24], thus giving a dry matter content of 91.02%. This suggests a nutrient-dense food material that can actually be utilized in many ways such as in-feed supplementation [3]. "The Ash content of the sample could be related to the mineral content and these minerals, which are mostly in form of chemical compounds, play numerous functions in the improvement of health in the body of organisms" [25]. The fat present in the seed is relatively high, in agreement with the result of Ifesan et al., [26], and may be an indication that *Persea americana* could be a good source of oil. The value falls within the range of values for most oil-producing seeds like soybeans [27]. "Generally, fats have many functions aside from insulation and conservation of body temperature in organisms, their fatty acid components such as lauric acid and polyunsaturated fatty acids have been reported to improve health" [7]. From this study, the crude fiber content in the seed of *P. americana* is high when compared with its content in the African pear seed, 3.17%, [3]. Adequate dietary fiber intake alters the colonic environment in such a way as to protect against colorectal diseases, lowering of serum cholesterol, and prevent hypertension, diabetes, constipation, and heart diseases [28]. "It provides protection by increasing fecal bulk, which dilutes the increased colonic bile acid concentrations that occur with a high-fat diet" [29]. In comparing the protein content in this study with other studies, the value agrees with the report of Ifesan et al., [26]. "Aside from contributing to diets, the

relative impact of proteins in the body system should not be overlooked. As chemical compounds, they repair and replace worn-out cells, form structural and globular materials that hold the body, form blood proteins and boost the immune system" [25]. In this study for proximate composition, the carbohydrate content is highest which is similar to the results of other studies as in Onuegbu et al., [3] and Ifesan et al., [26]. The carbohydrate content of the sample is related to energy generation [25]. Observed carbohydrates in the sample may be an indication that the sample could produce energy to power the cells and tissues of the body on consumption.

"Phytochemicals are important chemicals found virtually in plants and they are in different parts and at different concentrations" [30]. The phytochemical profile of *P. americana* seeds revealed that they are rich in phenols, flavonoids, and terpenoids (Fig. 2). Phenols have been extensively researched as disease preventives [30]. "Phenols detected in seeds of *P. americana* investigated in this study could further indicate their ability to act as anti-inflammatory, anticlotting, antioxidants, immune enhancers, anticarcinogenic and antiaging" [10]. The results for flavonoid and tannin contents were similar to those of Nwaoguikpe and Braide, [8]. "Flavonoids are potent water-soluble super antioxidants and free radical scavengers. They prevent oxidative cell damage, have strong anticancer activity, and protect against all stages of carcinogenesis" [31]. "In the intestinal tract, they lower the risk of heart disease, and inflammation and represent the most common and widely distributed groups of plant phenolic compounds" [7].

"Alkaloids are therapeutically significant plant secondary metabolites. The isolated pure form of alkaloids and their synthetic derivatives are used as basic medicinal agents for their analgesic and bactericidal effects" [32]. The presence of alkaloids in the seed extract could serve as a reason for its use in ethnomedicine [33]. "The content of tannins in *P. americana* may be responsible for their free astringency and bitter taste" [34]. They hasten the healing of wounds and inflamed mucous membrane [35] hence their use in the treatment of injuries. Tannins are known to bind irreversibly to proteins by forming insoluble complexes with them and thus rendering them indigestible by intestinal enzymes thereby interfering with their bioavailability [36]. The Saponin level in the seed sample was moderate and thus could not produce adverse

effects on the growth of animals [10]. “Some of the general characteristics of saponins include the formation of foams in an aqueous solution, hemolytic activity, and cholesterol-binding properties” [37]. The oxalate level in the seed sample was below the lethal dosage of soluble oxalate intake (3 – 5 g) [38], hence will have little or no adverse effect. The phytic acid content of the avocado raw seed is below the recommended daily allowance dose of 25mg/100g [39] hence will have a little anti-nutritional effect. “The anti-nutritional nature of phytic acid lies in its ability to chelate divalent minerals such as iron, calcium, copper, and zinc, rendering them biologically unavailable. Processing techniques (soaking and boiling) significantly ( $p \leq 0.05$ ) reduced the phytic acid content of avocado seeds” [40]. Phytic acids are the principal form of phosphorus in many seeds; 60–90% of phosphorus in seeds is present as phytic acid [41] hence lowering the total phosphorus value in avocados.

From this analysis, avocado seeds were found to be rich in beta-carotene and lycopene. Lycopene is a good antioxidant and a precursor for beta

carotene while beta carotene serves as a food colorant (responsible for the yellowish color of the seeds) and a precursor for vitamin A [42]. “This effect of RA is mediated mainly by retinoic acid receptors and varies among cell types. In mammary carcinoma cells, the retinoic acid receptor was shown to trigger growth inhibition by inducing cell cycle arrest, apoptosis, or both” [43]. Lycopene is a cellular inhibitor for various inflammatory processes.

Also, the result revealed that the seed sample contain a good amount of total phytosterols, stigmasterols, and campesterol. “The roles of most plant sterols have been documented. One strategy adopted in the reduction of cholesterol levels involves the use of plant sterols such as sitosterol, stigmasterol, and campesterol, whose mechanism of action despite their structural similarity to cholesterol involves the inhibition of cholesterol absorption by the intestinal cells” [44]. “The cholesterol-lowering potential of phytosterols is supposed to be caused by an inhibition of cholesterol absorption resulting from the higher solubility of phytosterols than of cholesterol in bile salt micelles” [9].





Fig. 2. Bar chart of phytochemical composition of *Persea americana* seed



**Fig. 3. Bar chart of the mineral composition of *Persea americana* seed**

The mineral composition result (Fig. 3) clearly indicates that *P. americana* seeds are good sources of macro minerals like phosphorus, potassium, calcium, and magnesium, except relatively low sodium. The concentrations of these minerals are not the same as reported in other studies, [10] and [25]. The varying composition reported by various studies could vary with season, environment, and condition or time evaluation. Phosphorus, calcium, and magnesium are required for the formation of bones and teeth, formation of the blood clot, the formation of cyclic AMP, and other second messengers, for body mechanisms [25]. Potassium is necessary for electrolyte balance and control of high pressure. This could also be implicated in the use of *P. americana* seed to treat high blood pressure in traditional medicine [10]. The low sodium level is an indication that the seed sample can not jeopardize blood pressure since the elevated level of sodium salt in the blood has been associated with high blood pressure in the body [25], but this may not be possible in a situation of higher potassium content. Trace elements such as molybdenum, and selenium, are needed in small quantities by the body; therefore concentrations of these elements found in *P. americana* seeds are of great nutritional importance [2]. Zinc, which is the

most prevalent trace element in this study, is essential for the production of insulin and hence can play a valuable role in the management of diabetes [2]. Zinc plays role in wound healing, iron is known for haem formation, and manganese and copper aid iron absorption in the body [25]. Selenium plays an important role in the antioxidant system by acting as a cofactor for glutathione peroxidase, enhancing alpha-tocopherol activities, and helping in the DNA repair mechanism. Copper is essential for hemoglobin synthesis, normal bone formation, and the maintenance of myelin within the nervous system [45,46].

## 5. CONCLUSION

*P. americana* seeds have appreciable nutrients in the form of carbohydrates, crude protein, crude fat, and fiber, hence they can be added to our dietary plan (such as homemade smoothies). The appreciable antioxidant activities exhibited by the seed extract are attributed to the presence of the phytochemicals such as flavonoids, phenols, lycopene, and the like. The total phytosterols present in *P. americana* seed could also serve as preventive measures against cardiovascular diseases since phytosterols exhibit hypocholesterolemia activities. These

possible dietary and drug functions of methanol seed extract of *P. americana*, have revealed the background for phytomedicinal usage of avocado pear seeds thereby demanding further studies to determine whether these seeds have possible usage in medicine and food industries.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. João JL, Érika HB, Rossana AC, Raimunda SN, José JC, Luciana MB, Selene MM, Marcos GR. Chemical composition, toxicity and larvicidal and antifungal activities of *Persea americana* (avocado) seed extracts. *Revista da Sociedade Brasileira de Medicina Tropical*. 2009;42(2):110-113.
2. Ajayi IA, Adesanwo O. Comparative study of the mineral element and fatty acid composition of *Dacryodes edulis* pulp and seed. *World Journal of Agricultural Sciences*. 2009;5:279-283.
3. Onuegbu NC, Nwuka MU, Ojukwu M, Kabu NO. Nutritional properties of African pear seed and performance of defatted cake in poultry feed formulations. *Animal Nutrition*. 2016;1(29):1-3.
4. Ortiz MA, Dorantes AL, Gallindez MJ, Cárdenas SE. "Effect of a novel oil extraction method on avocado (*Persea americana* Mill) pulp microstructure." *Plant Foods for Human Nutrition*. 2004;59(1): 11–14.
5. Ferrari RA, Colussi F, Ayub RA. Characterization of by-products of passion fruit industrialization. *Revista Brasileira de Fruticultura*. 2004;26:101-102.
6. Anaka ON, Ozolua RI, Okpo SO. Effect of the aqueous seed extract of *Persea americana* Mill (*Lauraceae*) on the blood pressure of sprague dawley rats. *African Journal Pharmacy and Pharmacology*. 2009;3(10):485-490.
7. Imafidon KE, Amaechina FC. Effects of aqueous seed extract of *Persea americana* Mill. (Avocado) on blood pressure and lipid profile in hypertensive rats. *Advance Biological Researches*. 2010;4(2):116-121.
8. Nwaoguikpe RN, Braide W. The effect of aqueous seed extract of *Persea americana* (avocado pear) on serum lipid and cholesterol levels in rabbits. *African Journal of Pharmacy and Pharmacology Research*. 2011;1(2):023-029.
9. Hirota Y, Nagao T, Watanabe Y, Suenaga M, Nakai S, Kitano M, Sugihara A, Shimada Y. Purification of steryl esters from soybean oil deodorizer distillate. *Journal of the American Oil Chemists' Society*. 2003;80:341-346.
10. Arukwe U, Amadi BA, Duru MKC, Agomuo EN, Adindu EA, Odika PC, Lele KC, Egejuru L, Anudike J. Chemical composition of *Persea americana* leaf, fruit and seed. *International Journal of Research and Reviews in Applied Sciences*. 2012;11(2):2-10.
11. Deepti D, Rachel MS, Gregory RZ, Joshua DL. Avocado (*Persea americana*) seed as a source of bioactive phytochemicals. *Current Pharmaceutical Design*. 2013;19:6133-6140.
12. Kumar AK, Ramachandra SS, Narsu L. Pharmacognostic and phytochemical investigation of roots of *Hibiscus micranthus* Linn. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 2010;1(4):324-337.
13. AOAC. Official method of proximate analysis of food (17<sup>th</sup> ed) Association of Official Analytical Chemists, Arlington, Virginia, U.S.A; 1999.
14. Merrill AL, Watt BK. Energy value of foods: Basis and derivation. *Agriculture Handbook No. 74*. Washington, DC, ARS United State Department of Agriculture; 1973.
15. Trease GE, Evans WC. *Phytochemicals. Pharmacognosy*. 13th ed. ELBS/Bailliere Tindal, London. 1989:345–346.
16. Harborne JB. *Phytochemical methods. A guide to modern techniques of plants analysis*. 2<sup>nd</sup> Edition, Chapman and Hall, Ltd., London, UK. 1973:149-188.
17. Young SM, Greaves JS. Influence of variety and treatment on phytic acid content of wheat. *Journal of Food Research*. 1940;5:103-105.
18. Osagie AU. Antinutritional factors in nutritional quality of plant foods. In "Adeniyi SA, Orjiekwe CL, Ehiagbonare JE. Determination of alkaloids and oxalates in some selected food samples in Nigeria. *African Journal of Biotechnology*. 1998;8(1):110-112.
19. Narayan S, Amit RS, Bishun DP, Kanti S. Investigation on phytochemical, antimicrobial activity and essential oil constituents of *Nardostachys jatamansi*

- DC. In different regions of Nepal. Journal of Coastal Life Medicine. 2016;4(1):56-60.
20. Pearson D. The chemical analysis of foods. Churchill Livingstone, Edinburgh. 6<sup>th</sup> Edition. 1974: 457.
  21. Barros L, Ferreira MJ, Queiros B, Ferreira ICFR, Batista P. Total phenols, ascorbic acid,  $\alpha$ -carotene and lycopene in portuguese wild edible mushrooms and their antioxidant activities. Food Chemistry. 2007;103:413-419.
  22. Larissa BDCA, Sarah LS, Marcos AMG, Magda RAF, Evani LA, Karina PR, Luiz AL. Total phytosterol content in drug materials and extracts from roots of *Acanthospermum hispidum* by UV-VIS spectrophotometry. Brazilian Journal of Pharmacognosy. 2013;23:736-742.
  23. American Public Health Association (APHA). Standard methods for the examination of water and waste water, 19<sup>th</sup> edition APHA, Washington, D.C. 1995;1644.
  24. Chikezie PC, Agomuo EN, Amadi BA. Proximate analysis In: Biochemistry Practical/Research method. A Fundamental Approach. Megasoft Publishers, Owerri. 2008;2:8-21.
  25. Olusanya JO. Proteins, In: Essentials of food and nutrition. Apex Books Limited, Lagos. 2008:13-21.
  26. Ifesan BOT, Olorunsola BO, Ifesan BT. Nutritional composition and acceptability of candy from avocado seed (*Persea americana*). International Journal of Agriculture Innovations and Researches. 2015;3(6):2319-1473.
  27. Krogdahl A, Bakk-Mckellep AM. Physiological mechanism in the impairment of nutrient digestion and transport capacity in salmon intestine. Soybean Association Hudson Iowa, USA. 2002:128.
  28. Ishida H, Suzunoh S, Sugiyana N, Innami S, Todoro T, Mackawa A. Nutritional evaluation of chemical components of leaves, stalk and stem of sweet potatoes (*Ipomea batatas*). Food Chemistry. 2000;68:359-367.
  29. Dillard CJ, German EM. Coconut: In support of good health in the 21st century. 1990:127.
  30. Duke J. Handbook of biological active phytochemicals and their activities. CRC Press, Boca Raton (Florida). 1992:99-131.
  31. Salah W, Miller NJ, Pangauga T, Bolwell GP, Rice E, Evans C. Polyphenolic flavonols as scavengers of aqueous phase radicals as chainbreaking antioxidant. Archives of Biochemistry and Biophysics. 1995;2:339-346.
  32. Stray F. The national guide to medicinal herbs and plants. Tiger Books International, London. 1988:1246.
  33. Ramos MR, Jerz G, Villanueva S, Lopez-Dellamary F, Waibe R, Winterhalter P. Two glucosylated abscisic acid derivatives from avocado seeds (*Persea americana* Mill. Lauraceae cv. Hass). Phytochemistry. 2004;65:955-962.
  34. Okwu DE. Phytochemicals and vitamin content of indigenous spices of South Eastern Nigeria. Journal Of Sustainable Agriculture and Environment. 2004;6:30-34.
  35. Okwu DE, Okwu ME. Chemical composition of spondiasombia Linn plant parts. Journal of Sustainable Agriculture and the Environment. 2004;6:140-147.
  36. Liener IE. Implications of antinutritional components in soybean foods. Critical Reviews in Food Science and Nutrition. 1994;34(1):31-67.
  37. Sodipo OA, Akiniyi J. Studies on certain characteristics of extracts from bark of *Pansinystaliamacruceras* (K. schum). Pierre Exbeille. Global Journal of Pure and Applied Science. 2000;6:83-87.
  38. Antia BS, Akpan EJ, Okon PA, Umoren IU. Nutritive and anti-nutritive evaluation of sweet potatoes leaves. Pakistan Journal of Nutrition. 2006;5(2):166-170.
  39. Hurrell RF, Jullert MB, Reddy SR, Lynch SD, Cook JD. Changes in anti-nutrients contents of edible vegetables under varied temperature and heating time. American Journal of Clinical Nutrition. 1992;56:573.
  40. Justina YT, Olukemi AO, Ajayi OO, Adegoke GO. Nutritional and antinutritional compositions of processed avocado (*Persea americana* Mill) seeds. Journal of Plant Science and Research. 2016;6(2):6-12.
  41. Mohammad A, Koji Y, Toshiki M, Yomi N, Katsumi S, Hiroaki S, Yoshifumi T. Superoxide anion radical scavenging activities of herbs and pastures in northern Japan determined using electron spin resonance spectrometry. International Journal of Biological Sciences. 2007 3:349-355.
  42. Susan DA. "Vitamin A", kirk-othmer encyclopedia of chemical technology (45). New York: John Wiley. 1998:99-107.



43. Niizuma H, Nakamura Y, Ozaki T, Nakanishi H, Ohira M. Bcl-2 is a key regulator for the retinoic acid-induced apoptotic cell death in neuroblastoma. *Oncogene*. 2006;25:5046-5055.
44. Evans WC. Traditional plant medicines as a source of new drugs. In Trease and Evans *Pharmacognosy*, 15<sup>th</sup> edition. Publishers, Elsevier. 2005:125-134.
45. Passmore R, Eastwood MA. In Davison Passmore, R., *Human Nutrition and Dietetics*, Churchill Livingstone, London. 1986:124-126.
46. Fite B. *The Healing Miracle of Coconut Oil*. Piccadilly Books Ltd, Healthwise Publications Colorado Springs, Co. 2000; 1-4.

---

© 2023 Omorevbarhia et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/97536>