



Antimicrobial Activity of Natural Intracanal Medicaments in Endodontic Management of Permanent Teeth: A Systematic Review of *In-vivo* Studies

**Subikshaa ^{a++*}, Rathna Piriyaanga ^{b,c#}, Geeth Deepika ^{b#},
Nivashini GSV ^{b†} and Anand Sherwood ^{b‡}**

^a CSI College of Dental Sciences and Research, Madurai, Tamil Nadu, India.

^b Department of Conservative Dentistry and Endodontics, CSI College of Dental Sciences and Research, Madurai, Tamil Nadu, India.

^c Department of Conservative Dentistry and Endodontics, Saveetha Dental College, Chennai, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration among all authors. Authors Subikshaa and AS has done the design of the study. Authors Subikshaa and RP has done the data collection. Authors Subikshaa, RP and GD has done the data extraction. Authors Subikshaa, RP, GD and NGSV has done the data analysis and authors Subikshaa, RP and GD contributed for preparation of manuscript. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/cjast/2024/v43i114439>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125486>

Systematic Review Article

Received: 20/08/2024

Accepted: 22/10/2024

Published: 26/10/2024

⁺⁺Intern;

[#]Reader;

[¶]PhD Scholar;

[†]Senior Lecturer;

[‡]Professor and Head;

^{*}Corresponding author: E-mail: subineelu3@gmail.com;

Cite as: Subikshaa, Rathna Piriyaanga, Geeth Deepika, Nivashini GSV, and Anand Sherwood. 2024. "Antimicrobial Activity of Natural Intracanal Medicaments in Endodontic Management of Permanent Teeth: A Systematic Review of In-Vivo Studies". *Current Journal of Applied Science and Technology* 43 (11):1-12. <https://doi.org/10.9734/cjast/2024/v43i114439>.

ABSTRACT

Background: Endodontic therapy has long been recognized as the modality for the elimination of microbial infections from the root canal system. Traditionally, calcium hydroxide has been prevalent in the history of intracanal medicaments because of their inherent antimicrobial activity. However, interest in recent years has been in the area of endodontic therapy incorporating natural intracanal medicaments from plant extracts into their protocols. The biocompatibility of these agents, its reduced cytotoxic potential, and other therapeutic properties such as anti-inflammatory and tissue regenerative have all been promising. But the clinical effectiveness of these natural agents is yet to be clearly established, especially in permanent teeth.

Objective: This systematic review aims at assessing the effectiveness of natural intracanal medicaments in the endodontic treatment of permanent teeth by pooling evidence from *in vivo* studies regarding their antimicrobial activity, as compared to traditional synthetic agents.

Methodology: Electronical databases, such as Google Scholar, Web of Science, PubMed, and Scopus were searched for articles published between 2014 and 2021. Rayyan software was used in the screening and management of all articles and, during the study selection, title and abstract screening were done independently by two reviewers. Meta-analysis assessed the pooled effect size of such agents.

Results: Of nearly 2500 articles, only 6 qualified for inclusion. The natural agents Neem, Curcumin, and Propolis demonstrated high antimicrobial activity compared to conventional agents such as calcium hydroxide and chlorhexidine. Neem had the highest antibacterial activity, and Curcumin demonstrated anti-inflammatory effects superior to all else. While results from these natural medicaments demonstrated a broad efficacy, significant heterogeneity was found among the studies at 96% ($I^2 =$).

Conclusion: Natural intracanal medicaments are a promising future of efficacy: as good as or even better than traditional synthetic agents applied in endodontic therapy. Some of them are Aloe vera, Curcumin, Neem, and Propolis, which have been shown to exhibit not only good antimicrobial efficacy but also some degree of anti-inflammatory and tissue-regenerating properties. However, more standardized clinical trials should be designed and well executed to validate their effectiveness, thus defining clear clinical guidelines for permanent teeth use.

Keywords: Antimicrobial activity; antibacterial effects; endodontics; intracanal medicaments; natural materials.

1. INTRODUCTION

Endodontic therapy is crucial for managing microbial infections within the root canal system, necessitating the use of effective intracanal medicaments to eradicate pathogens and promote periapical healing (Athanassiadis, et al. 2007). Conventional agents, such as calcium hydroxide, are widely used in endodontic treatment due to their potent antimicrobial properties and ability to stimulate the formation of mineralized tissue. Chlorhexidine (CHX) is also a key agent, particularly valued for its efficacy in retreatment cases (Digole et al., 2020). Triple antibiotic paste (TAP), composed of ciprofloxacin, metronidazole, and minocycline, has gained prominence as an effective intracanal medicament, especially in cases involving regenerative endodontics [Ordinola-Zapata, et al. 2022].

Natural intracanal medicaments, derived from plant extracts, present a compelling option due to

their biocompatibility, reduced cytotoxicity, and additional therapeutic benefits. Many of these substances exhibit inherent antimicrobial properties while also offering anti-inflammatory and tissue-regenerative effects, making them attractive candidates for endodontic therapy (Karobari et al. 2014). The ability of these agents to mitigate inflammation and enhance healing processes aligns with the evolving focus on minimally invasive and biologically compatible treatments in dentistry (Tirukkolluru & Thakur, 2019).

Despite the promising potential of natural intracanal medicaments, uncertainty remains regarding their clinical efficacy, particularly in permanent teeth (Al-Sabawi et al., 2020). While numerous *in vitro* studies have demonstrated the antimicrobial activity of various natural substances, their clinical relevance is often limited. Therefore, understanding their effectiveness *in vivo* is essential for optimizing treatment outcomes and advancing evidence-


based endodontic practice (Lillygrace et al., 2021). Randomized controlled trials and prospective clinical studies are imperative to thoroughly evaluate the true effectiveness of these natural agents in clinical settings, establishing standardized protocols for their application in routine endodontic therapy (Rathee et al., 2020).

This systematic review and meta-analysis aim to consolidate existing literature on the antibacterial efficacy of natural intracanal medicaments, highlighting their mechanisms of action and potential advantages over conventional agents. By synthesizing the findings of these studies, we aim to provide clinicians with valuable insights

into the clinical effectiveness of natural intracanal medicaments and emphasize the importance of conducting well-designed, standardized research to inform evidence-based endodontic practice.

2. MATERIALS AND METHODS

This systematic review followed PICO statement for including the studies, on patients undergoing endodontic treatment in permanent teeth. The intervention under investigation is the utilization of natural materials as intracanal medicaments during endodontic treatment. These natural materials encompass a range of substances including propolis, neem, garlic extract, and others derived from plants.



In total 6 articles are selected based on inclusion and exclusion criteria.

Fig. 1. PRISMA flowchart for included studies

These materials are applied with the aim of disinfecting the root canal system and facilitating healing. The comparison group comprises conventional synthetic medicaments commonly employed in endodontic treatment. The primary outcome of interest is the assessment of the antimicrobial properties of the intracanal medicaments. This assessment is based on the reduction in microbial colony counts within the root canal system.

2.1 Search Strategy

A comprehensive search strategy was employed, utilizing various electronic databases including PubMed, Scopus, Web of Science, and Google Scholar. The search terms utilized encompassed keywords intracanal medicament OR root canal medicament OR endodontic medicament OR root canal disinfectant AND natural materials OR plant-based materials OR organic materials OR non-synthetic materials AND antimicrobial efficacy OR antimicrobial activity OR antibacterial effectiveness OR microbial eradication OR microbial suppression. The Rayyan software was employed for efficient screening and management of search results.

2.2 Study Selection

The study selection followed Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) guidelines. Inclusion criteria for study selection encompassed studies published between 2017 to 2021, *in vivo* studies conducted exclusively in permanent teeth, evaluating the efficacy of natural materials as intracanal medicaments in endodontic treatment, and reporting microbial colony counts as an outcome measure. Exclusion criteria included studies conducted in primary teeth and those not meeting the specified inclusion criteria (Fig. 1).

2.3 Data Extraction

Data extraction involved gathering pertinent information from each included study, including author details, publication year, study design, sample size, the natural material used as an intracanal medicament, and outcomes related to microbial colony counts (Table 1).

2.4 Quality Assessment

The quality assessment of included studies was conducted using the Risk of Bias 2.0 (ROB 2.0) tool. This comprehensive tool evaluates various

domains of study quality, including randomization procedures, allocation concealment, blinding of participants and personnel, completeness of outcome data, selective outcome reporting, and other sources of bias. Any disagreements between reviewers (1st author and 2nd author) were handled through discussion or consultation with a third reviewer (3rd Author) to reach a consensus.

3. RESULTS

3.1 Quality Assessment Results

The quality assessment using the ROB 2.0 tool found that the majority of studies had a low risk of bias. Overall, while the majority of the included studies demonstrated a high level of methodological rigor, Samta Khetarpal et al. (2014) stood out as having a high risk of bias in certain aspects (Fig. 2).

3.2 Data Extraction Result

A total of six studies were included in the systematic review, comprising research conducted in India and Saudi Arabia between 2014 and 2021. The studies varied in design, with two randomized controlled trials, one non-randomized controlled trial, one experimental study, one prospective randomized controlled study, and one randomized single-blinded clinical trial.

Regarding the antimicrobial efficacy of different intracanal medicaments, including natural materials and conventional synthetic chemicals, notable findings emerged as mentioned in Table 1. Overall, the reviewed studies provide valuable insights into the antimicrobial efficacy of various intracanal medicaments, emphasizing the potential of natural materials as alternatives to conventional synthetic chemicals in root canal therapy.

3.3 Meta Analysis Result

In the meta-analysis, the effect sizes varied across the six individual studies. The pooled effect size, represented by the diamond at the bottom of the forest plot, indicated a Inverse variance method to compare the standardized mean difference (SMD), there is no statistical difference between the two cohorts, the summarized standardized mean difference (SMD) is -0.9 with a 95% confidence interval of -11.47 - 9.68. The test for overall effect does not

show a significant effect. A significant heterogeneity was detected ($p < 0.01$), suggesting inconsistent effects in magnitude and/or direction. The I^2 value indicates that 96% of the variability among studies arises from heterogeneity rather than random chance.

These findings highlight the significant impact of the experimental condition compared to the control, warranting further investigation and consideration of potential sources of heterogeneity in future research (Fig. 3).




Fig. 2. Risk of Bias assessment graph using ROB 2.0 tool




Fig. 3. Meta analysis results shown as forest plot mentioning all included studies

Table 1. Data extracted results from the included studies

Name year Region	Sample size	Group	Study design	Independent variables	Dependent variables	Outcome measures	Results	Main findings	Limitations	Summary
Samta Khetarpal 2014 India [4]	Total: 33	Group A – Neem Group B – Curcumin Group C – Aloe vera Control - Chlorhexidine	Non-randomized controlled trial with a parallel design	Type of intracanal medicament used (Neem, Curcumin, Aloe vera, Chlorhexidine)	Antimicrobial and anti-inflammatory properties of the intracanal medicaments (Neem, Curcumin, Aloe vera)	The outcome measured in the study includes antimicrobial activity assessed through microbial colony count scores, anti-inflammatory activity assessed through the Visual Analogue Pain Scale, and the comparison of antimicrobial and anti-inflammatory properties between the test groups (Neem, Curcumin, Aloe vera) and the control group (Chlorhexidine).	The results were statistically non-significant when Neem was compared with Chlorhexidine, and statistically significant when Curcumin and Aloe vera were compared with Chlorhexidine in terms of anti-microbial effect.	Neem showed the highest antibacterial activity, while Curcumin exhibited the highest anti-inflammatory activity compared to Aloe vera and Neem. Curcumin and Aloe vera showed statistically significant results compared to Chlorhexidine in terms of antimicrobial and anti-inflammatory properties.	Many limitations include no hypothesis is stated, inclusion and exclusion criteria is not given properly, whether the guidelines are followed are not given properly.	The study evaluated the antimicrobial and anti-inflammatory properties of Neem, Curcumin, and Aloe vera as intracanal medicaments in root canal therapy, highlighting their effectiveness compared to Chlorhexidine.
Chandrasekar 2020 India [5]	Total: 45	Group I: Propolis with moxifloxacin Group II: TAP Group III: Calcium hydroxide	Randomized clinical trial, controlled	Type of intracanal medicament used (Propolis with moxifloxacin, TAP, Calcium hydroxide),	Viable colony-forming unit counts of <i>E. faecalis</i> and <i>Streptococcus</i> spp. in the root canals after instrumentation and medication sessions	Reduction in microbial count of <i>Streptococcus</i> spp. and <i>Enterococcus faecalis</i> in the root canals of Type II DM patients with chronic apical periodontitis after treatment with different intracanal medicaments	At the end of the study, intragroup differences in <i>E. faecalis</i> and <i>Streptococcus</i> species mean CFU counts in postinstrumentation samples (S1) and postmedication samples (S2) were statistically significant for each of the Groups 1, II, and III. However, intergroup differences and pair wise comparison of <i>E. faecalis</i> and <i>Streptococcus</i> species mean CFU counts from S1 to S2 were statistically nonsignificant ($P > 0.05$) among all the study groups.	- The study showed a significant reduction in microbial count in all study groups from the first to the second session, with no significant difference in the decrease of microbial load between the three groups at the end of 1 week. - The antimicrobial efficacy of TAP, propolis with moxifloxacin, and calcium hydroxide was concluded to be comparable.	The limitations of the study include issues related to the sensitivity and specificity of the culture techniques and CFU counts, as well as the experimental design.	The paper discusses the comparable antimicrobial efficacy of TAP, propolis with moxifloxacin, and calcium hydroxide in diabetic patients with chronic apical periodontitis, emphasizing the importance of intracanal medicaments in root canal treatment success.
Goldy Rathee 2021 India [6]	Total: 30	Group I: 2% Chlorhexidine gluconate. Group II: Neem juice extract	Randomized controlled trial	Types of intracanal medicaments used (2% Chlorhexidine	The antibacterial efficacy measured by counting	Reduction in colony-forming units (CFU) at	The results showed a statistically	The main findings of the study suggest that herbal	The limitations of the study are not explicitly stated in	The paper evaluates the antimicrobial

Name year Region	Sample size	Group	Study design	Independent variables	Dependent variables	Outcome measures	Results	Main findings	Limitations	Summary
		Group III: Tulsi juice extract		gluconate, Neem juice extract, and Tulsi juice extract)	colony-forming units from the samples taken at different stages	different stages of root canal treatment: after access opening (S1), after biomechanical preparation and irrigation (S2), and after placing intracanal medicaments (S3)	significant reduction among all the groups during intragroup comparison. Among these stages, a significant reduction was seen in group I and group II, group I and group III.	products exhibit significant antimicrobial activity in treating primary endodontic infections compared to 2% CHX, supporting their recommendation as effective endodontic irrigants and medicaments. The study underscores the potential of herbal alternatives in root canal therapy.	the paper. The only indirect reference to limitations is the mention of "within the limitations of this study" before drawing the conclusion. The paper does not provide a specific list of limitations such as sample size, generalizability, bias, or methodological errors.	efficacy of herbal products as root canal irrigants and medicaments, showing their significant activity compared to chlorhexidine gluconate, supporting their recommendation for endodontic use.
NawalA Al-Sabawi 2020 Saudi Arabia [7]	Total - 60	Group I: NS (control -ve) Group II: 2% CHX (control +ve) Group III: FX Group IV: 8% FL Group V: 8% OL Group VI: 8% SP	Experimental study without randomization, blinding, control groups, or placebos, involving both <i>in vitro</i> and <i>in vivo</i> experiments.	- Concentrations of plant extracts (2%, 4%, and 8%) - Types of plant extracts (FX, FL, OL, SP) - Groups for <i>in vivo</i> study (NS, 2% CHX, FX, 8% FL, 8% OL, 8% SP)	The dependent variables in Nawal A Al-Sabawi, Ali M Al-Naimi, Eman M Yahya (2020) are: - The antibacterial effect of different plant extracts (Ficus carica latex, Ficus carica leaves, olive leaves, and <i>Salvadora persica</i> roots) and 2% chlorhexidine against <i>Enterococcus faecalis</i> when used as intracanal medicaments - The mean absorbance values of the different plant extracts and chlorhexidine <i>in vitro</i> - The bacterial counts of the different plant extracts and chlorhexidine <i>in vivo</i>	Antibacterial effect of plant extracts and chlorhexidine against <i>Enterococcus faecalis</i> for the elimination of <i>E. faecalis</i> from the root canal when used as intracanal medicaments	The study found that <i>Ficus carica</i> latex, <i>Ficus carica</i> leaves, olive leaves, and <i>Salvadora persica</i> can be used successfully as intracanal medicaments due to their antibacterial effect against most microbial flora in the root canal, especially <i>E. faecalis</i> . The <i>in vivo</i> study confirmed their antibacterial effect against <i>E. faecalis</i> , which was considered the first clinical study evaluating the effectiveness of these intracanal medicaments against <i>E. faecalis</i> . These plant extracts were biocompatible and some of them had chelating effect. The bacterial counts of these plant extracts were nearly similar and	Ficus carica latex and ethanolic extracts of <i>Ficus carica</i> leaves, olive leaves, and <i>Salvadora persica</i> roots were effective in eliminating <i>Enterococcus faecalis</i> , potentially serving as alternative intracanal medicaments. - The antibacterial effect of these plant extracts was comparable to 2% chlorhexidine. - These plant extracts could be valuable in endodontic treatments for disinfection of the root canal system.	sample size and distribution, potential bias in patient selection,	The paper evaluates the antibacterial activity of different plant extracts and chlorhexidine against <i>Enterococcus faecalis</i> , concluding that certain plant extracts can be effective intracanal medicaments.

Name year Region	Sample size	Group	Study design	Independent variables	Dependent variables	Outcome measures	Results	Main findings	Limitations	Summary
							significantly not different from each other, and significantly different from NS.			
Vasantha R Digole 2020 India [8]	Total: 45	Group I – Curcumin Group II – Aloe vera Group III - Calcium hydroxide	prospective randomized controlled study	Type of intracanal medicament used (Curcumin, Aloe vera, Calcium hydroxide)	Post-operative bacterial CFU counts	Percentage decrease in bacterial numbers in postoperative samples compared to preoperative samples for each group	The study analyzed the effects of disinfection procedures on microbial colony counts in endodontic treatments using Curcumin, Aloe vera, and Calcium hydroxide. The data was statistically analyzed using Statistical Package for Social Sciences (SPSS). P-value<0.05 is considered to be statistically significant.	Curcumin showed the highest antibacterial efficacy as an intracanal medicament, followed by calcium hydroxide and aloe vera.	The limitations of the study include a small sample size per group, inaccessibility of paper points to accessory canals, and the suggestion for a longer follow-up study and radiographic assessment of periapical lesions.	The paper evaluates the antimicrobial efficacy of curcumin, aloe vera, and calcium hydroxide as intracanal medicaments, with curcumin showing the highest antibacterial efficacy.
Edara Lillygrace 2021 India [9]	Total: 30	Group I— triple antibiotic paste Group II - propolis	Randomized single-blinded clinical trial	Type of intracanal medicament used (Triple antibiotic paste and Propolis extract)	Bacterial count in the root canal samples collected at different stages of treatment	Bacterial colony count in the root canals at three different time intervals: after access opening, after irrigation, and after 3-4 weeks	Both group results are shown with their standard deviations, as well as p-values. The p-values for S1 in both groups were significant. However, none of the mean colony counts for the two groups were statistically significant, meaning there were no significant differences in antibacterial efficacy between the two groups.	propolis was found to be more effective at reducing bacterial colonies in the root canal system. The mean colony count of the samples treated with propolis was significantly lower than that of samples treated with triple antibiotic paste	small sample size, no long term follow-up (Not mentioned in article)	The study evaluates propolis and triple antibiotic paste as intracanal medicaments in young permanent teeth, finding propolis to have comparable antimicrobial efficacy to triple antibiotic paste.

4. DISCUSSION

The results of this systematic review and meta-analysis underscore the potential of natural intracanal medicaments as viable alternatives to conventional synthetic agents in endodontic therapy. Various natural agents, including *Aloe barbadensis* (Aloe vera), *Curcuma longa* (Curcumin), *Azadirachta indica* (Neem), *Ocimum sanctum* (Tulsi), *Propolis*, *Salvadora persica* (Miswak), and *Ficus carica* (Fig), were highlighted for their antimicrobial efficacy against endodontic pathogens.

4.1 Aloe vera (*A. vera*)

The antibacterial properties of *A. vera* are primarily due to its ability to disrupt the bacterial cell membrane, inhibit cellular respiration, and interfere with membrane synthesis, leading to cell leakage and subsequent bacterial death. *Aloe vera* (*Aloe barbadensis miller*) exhibits antimicrobial activity through its bioactive compounds like anthraquinones, saponins, and aloin, which disrupt microbial cell membranes and inhibit protein synthesis. It targets both Gram-positive and Gram-negative bacteria, offering anti-inflammatory and healing benefits [Radha and Laxmipriya, 2014], [Nejatzadeh-Barandozi, 2013]. [The gel contains active ingredients such as anthraquinones, tannins, myristic acid, curcumin, and nimbidin, which exhibit potent antibacterial and anti-adherence properties, thereby reducing bacterial colonization and adhesion to dentin [Monica and Monisha, 2014]. Aloe vera also stimulates fibroblast growth and collagen synthesis, promotes pulp cell proliferation, differentiation, and extracellular matrix mineralization, making it a potential agent for use as a medicament in regenerative cases [Songsiripraduboon et al. 2017, Carvalho et al. 2018, Lowther et al. 2012, Sholehva et al. 2016]. However, the inferior molecule-binding capacity of *A. vera* to *E. faecalis*, and limited membrane permeability, reduce its efficacy against this specific bacterium. Additionally, compared to other herbal remedies that have been tried, anthraquinones in *A. vera* might not be as effective against *E. faecalis* [Ghasemi et al. 2020].

4.2 Curcumin (*C. longa*)

Curcumin has shown moderate effectiveness in disinfecting dentinal tubules due to its ability to disrupt the bacterial extracellular polysaccharide matrix, affecting cell wall integrity [Yadav et al.

2018]. As a polyphenolic compound, curcumin strongly inhibits bacterial proliferation by hindering the dynamics of FtsZ assembly within the Z-ring, a critical component for bacterial cell division [Kaur et al. 2010]. Furthermore, photoactivated curcumin has been found to be as effective as triple antibiotic paste (TAP) and was able to penetrate deeper into the dentinal tubules [Devaraj et al. 2016, Tyagi et al. 2015]. However, more research is needed to identify the complete mechanism of action of curcumin as an antibacterial agent.

4.3 Neem (*Azadirachta indica*)

The antibacterial properties of neem are largely attributed to nimbidin, a bitter principle extracted from neem that exhibits disrupting microbial cell walls, increasing membrane permeability, and inhibiting protein synthesis [Siddiqui 1942]. Margolone, margolonone, and isomargolonone, tricyclic diterpenoids isolated from the stem bark, have also been shown to possess potent antibacterial activities [Pennington 1981].

4.4 Tulsi (*Ocimum sanctum*)

The antibacterial activity of Tulsi is due to the presence of linolenic acid, which has been shown to inhibit the growth of a wide range of bacteria [Singh et al. 1996]. Tulsi also possesses immunomodulatory effects, enhancing levels of interferon, interleukin-4, and T-helper cells, thereby strengthening the host's defense against infections [Mondal et al. 2011]. Moreover, the ability of Tulsi to reduce silver ions to silver nanoparticles adds to its antibacterial efficacy against both Gram-positive and Gram-negative bacteria [Singhal et al 2011].

4.5 Propolis

Propolis exerts its antibacterial effect by increasing the permeability of the bacterial cell membrane, disrupting membrane potential, and inhibiting ATP production, which decreases bacterial motility [Sforcin 2016, Kędzia 2013]. Its efficacy is generally higher against Gram-positive bacteria, due to the distinct structural differences in the outer membrane of Gram-negative bacteria and the presence of hydrolytic enzymes that degrade propolis' active components. Artepillin C, a phenolic compound found in Brazilian propolis, has been shown to exhibit strong antibacterial activity against MRSA *S. aureus* [Veiga et al. 2017].

4.6 *Salvadora persica* (Miswak)

The antibacterial efficacy of *S. persica* is mainly due to the presence of flavonoids in its ethanolic extract, which interact with bacterial cell walls and inhibit bacterial growth [Abdallah and Al-Harbi 2015]. Flavonoids exert multiple actions, such as inhibiting nucleic acid synthesis, energy metabolism, and disrupting cytoplasmic membrane function.

4.7 *Ficus carica* (Fig. leaves)

The antibacterial activity of fig. leaves is attributed to the presence of flavonoids, tannins, and terpenoids [Tchombe and Louajri, 2015]. Flavonoids interfere with bacterial energy metabolism and disrupt membrane integrity, causing bacterial cell death [Cushnie and Lamb 2005, Nazzaro et al. 2013]. Additionally, tannins bind to bacterial cell walls, leading to protein denaturation and disruption of bacterial metabolism [Dziąło et al. 2016]. The terpenoids present in fig. leaves may further disrupt bacterial membranes, enhancing their bactericidal effects.

Overall, the herbal agents reviewed in this study show varied mechanisms of antibacterial action, including cell membrane disruption, protein denaturation, and inhibition of bacterial metabolism. While some agents, like *C. longa* and *Propolis*, have shown superior penetration and deeper disinfecting abilities, others like *A. vera* and *Tulsi* provide additional benefits like anti-inflammatory effects and immune modulation. The observed variability in the efficacy of these natural medicaments can be attributed to differences in study design, sample sizes, concentrations used, and variations in intervention protocols.

From a clinical perspective, the transition from conventional agents to natural medicaments in endodontics offers several potential benefits, such as improved biocompatibility, reduced cytotoxicity, and additional therapeutic effects, including anti-inflammatory and tissue regenerative properties. However, the current evidence limited by the lack of standardized protocols and the limited number *in vivo* studies, making it challenging to establish definitive clinical guidelines for the use of these agents in permanent teeth. Future research should focus on addressing these limitations by employing well-designed randomized controlled trials with larger sample sizes and standardized methodologies.

5. CONCLUSION

This systematic review demonstrates the potential of natural intracanal medicaments as effective alternatives to conventional synthetic agents in endodontic therapy. The evidence supports the antimicrobial efficacy of various natural substances, including Aloe vera, Curcumin, Neem, and Propolis, which also offer additional benefits like anti-inflammatory and tissue-regenerative properties.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology.

Details of the AI usage are given below: Grammer correction and logical flow

1. Quillbot, Grammarly and ChatGPT 3

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abdallah, E., & Al-Harbi, K. (2015). Phytochemical screening and antibacterial activity of crude aqueous and ethanol extracts of *Salvadora persica* L. stem (Miswak) from Saudi Arabia. *Journal of Phytopharmacology*, 4, 243–247. <https://doi.org/10.31254/phyto.2015.4501>
- Al-Sabawi, N. A., Al-Naimi, A. M., & Yahya, E. M. (2020). An *in vitro* and *in vivo* antibacterial effect of different plant extracts on *Enterococcus faecalis* as intracanal medicament. *Journal of International Oral Health*, 12, 362–369. https://doi.org/10.4103/jioh.jioh_324_19
- Athanassiadis, B., P. V. Abbott, and Laurence J. (2007). Walsh. "The use of calcium hydroxide, antibiotics and biocides as antimicrobial medicaments in endodontics." *Australian dental journal* 52: S64-S82.
- Carvalho N. C., Guedes S. A. G., Albuquerque-Júnior R. L. C. et al. (2018). Analysis of

- Aloe vera cytotoxicity and genotoxicity associated with endodontic medication and laser photobiomodulation, *Journal of Photochemistry and Photobiology B: Biology*, 178, 348–354.
<https://doi.org/10.1016/j.jphotobiol.2017.11.027>, 2-s2.0-85035363520
- Cushnie, T. P., & Lamb, A. J. (2005). Antimicrobial activity of flavonoids. *International Journal of Antimicrobial Agents*, 26, 351–352.
- Devaraj, S., Jagannathan, N., & Neelakantan, P. (2016). Antibiofilm efficacy of photoactivated curcumin, triple and double antibiotic paste, 2% chlorhexidine, and calcium hydroxide against *Enterococcus faecalis* in vitro. *Scientific Reports*, 6, Article 24797.
<https://doi.org/10.1038/srep24797>
- Digole, V. R., Dua, P., Shergill, S. P. S., Pathak, P., Kumar, V., & Prakash, P. (2020). Comparative evaluation of antimicrobial efficacy of calcium hydroxide, curcumin, and aloe vera as an intracanal medicament: An *in vivo* study. *IP Indian Journal of Conservative and Endodontics*, 5(3), 114–119.
<https://doi.org/10.18231/ijce.2020.027>
- Działo, M., Mierziak, J., Korzun, U., Preisner, M., Szopa, J., & Kulm, A. (2016). The potential of plant phenolics in prevention and therapy of skin disorders. *International Journal of Molecular Sciences*, 17(1), Article 1.
<https://doi.org/10.3390/ijms17010001>
- Ghasemi, N., Behnezhad, M., Asgharzadeh, M., Zeinalzadeh, E., & Kafil, H. S. (2020). Antibacterial properties of aloe vera on intracanal medicaments against *Enterococcus faecalis* biofilm at different stages of development. *International Journal of Dentistry*, 2020, Article 8855277.
<https://doi.org/10.1155/2020/8855277>
- Karobari, M. I., Adil, A. H., Assiry, A. A., Basheer, S. N., Noorani, T. Y., Pawar, A. M., Marya, A., Messina, P., & Scardina, G. A. (2022). Herbal medications in endodontics and its application—A review of literature. *Materials*, 15(9), 3111.
<https://doi.org/10.3390/ma15093111>
- Kaur, S., Modi, N. H., Panda, D., & Roy, N. (2010). Probing the binding site of curcumin in *Escherichia coli* and *Bacillus subtilis* FtsZ—a structural insight to unveil antibacterial activity of curcumin. *European Journal of Medicinal Chemistry*, 45, 4209–4214.
- Kędzia, B., & Holderna-Kędzia, E. (2013). Aktywność antybiotyczna propolisu krajowego i europejskiego. *Postępy Fitoterapii*, 2, 97–107.
- Khetarpal, S., & Bansal, A. (2014). Comparison of anti-bacterial and anti-inflammatory properties of neem, curcumin, and aloe vera in conjunction with chlorhexidine as an intracanal medicament—An *in vivo* study. *DJAS*, 2(III), 130–137.
<https://doi.org/10.1055/s-0038-1671999>
- Lillygrace, E., Kethineni, B., Puppala, R., et al. (2021). Antimicrobial efficacy of triple antibiotic paste and propolis as an intracanal medicament in young permanent teeth: An *in vivo* study. *International Journal of Clinical Pediatric Dentistry*, 14(2), 243–248.
- Lowther W., Lorick K., Lawrence S. D., and Yeow W.-S. (2012). Expression of biologically active human interferon alpha 2 in Aloe vera, *Transgenic Research*. 21, no. 6, 1349–1357.
<https://doi.org/10.1007/s11248-012-9616-0>, 2-s2.0-84870385839
- Mondal, S., Varma, S., Bamola, V. D., Naik, S. N., Mirdha, B. R., Padhi, M. M., et al. (2011). Double-blinded randomized controlled trial for immunomodulatory effects of Tulsi (*Ocimum sanctum* Linn.) leaf extract on healthy volunteers. *Journal of Ethnopharmacology*, 136, 452–456.
- Monica, B., & Monisha, R. (2014). Aloe vera in dentistry—A review. *Journal of Dental and Medical Sciences*, 13, 18–22.
- Nazzaro, F., Fratianni, F., De Martino, L., Coppola, R., & De Feo, V. (2013). Effect of essential oils on pathogenic bacteria. *Pharmaceuticals*, 6, 1451–1474.
- Nejatzadeh-Barandozi, F. (2013). Antibacterial activities and antioxidant capacity of Aloe vera. *Organic Medicinal Chemistry Letters*, 3, 5. <https://doi.org/10.1186/2191-2858-3-5>
- Ordinola-Zapata, R., Noblett, W. C., Perez-Ron, A., Ye, Z., & Vera, J. (2022). Present status and future directions of intracanal medicaments. *International Endodontic Journal*, 55(Suppl 3), 613–636.
<https://doi.org/10.1111/iej.13731>
- Pennington, T. D. (1981). *Flora Neotropica*. New York Botanical Garden, NY Monograph No. 28.
- Radha, M. H., & Laxmipriya, N. P. (2014). Evaluation of biological properties and

- clinical effectiveness of Aloe vera: A systematic review. *Journal of Traditional and Complementary Medicine*, 5(1), 21–26.
<https://doi.org/10.1016/j.jtcme.2014.10.006>
- Rathee, G., Tandan, M., & Mittal, R. (2020). Evaluation of antimicrobial efficacy of commercially available herbal products as irrigants and medicaments in primary endodontic infections: *In vivo* study. *World Journal of Dentistry*, 11(6), 488–493.
- Sforcin, J. M. (2016). Biological properties and therapeutic applications of propolis. *Phytotherapy Research*, 30, 894–905.
<https://doi.org/10.1002/ptr.5605>
- Sholehvar F., Mehrabani D., Yaghmaei P., and Vahdati A. (2016). The effect of Aloe vera gel on viability of dental pulp stem cells. *Dental Traumatology*. 32(5), 390–396, <https://doi.org/10.1111/edt.12272>, 2-s2.0-84988890287.
- Siddiqui, S. (1942). *Curr. Science*, 11, 278–279.
- Singh, S., Majumdar, D. K., & Rehan, H. M. (1996). Evaluation of anti-inflammatory potential of fixed oil of *Ocimum sanctum* (Holy basil) and its possible mechanism of action. *Journal of Ethnopharmacology*, 54, 19–26.
- Singhal, G., Bhavesh, R., Kasariya, K., Sharma, A. R., & Singh, R. P. (2011). Biosynthesis of silver nanoparticles using *Ocimum sanctum* (Tulsi) leaf extract and screening its antimicrobial activity. *Journal of Nanoparticle Research*, 13, 2981–2988.
- Songsiripradubboon S., Kladkaew S., Trairatvorakul C. et al. (2017). Stimulation of dentin regeneration by using acemannan in teeth with lipopolysaccharide-induced pulp inflammation, *Journal of Endodontics*. 43(7), 1097–1103.
<https://doi.org/10.1016/j.joen.2017.01.037>, 2-s2.0-85018945638
- Tchombe, L. N., & Louajri, A. (2015). Therapeutic effects of *Ficus carica* leaves: A brief review. *ARNP Journal of Science and Technology*, 5, 37–41.
- Tirukkolluru, C., & Thakur, S. (2019). Comparative evaluation of triple antibiotic paste, propolis with moxifloxacin, and calcium hydroxide as intracanal medicaments against *Streptococcus spp.* and *Enterococcus faecalis* in type II diabetes mellitus patients: A randomized clinical trial. *Contemporary Clinical Dentistry*, 10, 191–196.
https://doi.org/10.4103/ccd.ccd_195_18
- Tyagi, P., Singh, M., Kumari, H., et al. (2015). Bactericidal activity of curcumin I is associated with damaging of bacterial membrane. *PLOS ONE*, 10(3), Article e0121313.<https://doi.org/10.1371/journal.pone.0121313>
- Veiga, R. S., De Mendonça, S., Mendes, P. B., Paulino, N., Mimica, M. J., Lagareiro Netto, A. A., & Marcucci, M. C. (2017). Artepillin C and phenolic compounds responsible for antimicrobial and antioxidant activity of green propolis and *Baccharis dracunculifolia* DC. *Journal of Applied Microbiology*, 122, 911–920.
<https://doi.org/10.1111/jam.13400>
- Yadav, R. K., Tikku, A. P., Chandra, A., et al. (2018). A comparative evaluation of the antimicrobial efficacy of calcium hydroxide, chlorhexidine gel, and a curcumin-based formulation against *Enterococcus faecalis*. *National Journal of Maxillofacial Surgery*, 9(1), 52–55.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/125486>