



Epidemiology and Management of MYMV Disease in Mung Bean (*Vigna radiata* (L.))

Preety Verma ^{a*}, Vinod Kumar Malik ^a, Pooja Sangwan ^a,
Rakesh Kumar ^a, Pankaj Yadav ^a, Mamta Khaiper ^b,
Bajrang Lal Sharma ^c, Niharika Sheoran ^a, Nisha Yadav ^a,
Monika Jangra ^b and Janvi Malik ^a

^a Department of Plant Pathology, College of Agriculture, CCSHAU, Hisar, 125004, India.

^b Department of Forestry, College of Agriculture, CCSHAU, Hisar, 125004, India.

^c Department of Entomology, College of Agriculture, CCSHAU, Hisar, 125004, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i31685

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97085>

Original Research Article

Received: 23/12/2022

Accepted: 28/02/2023

Published: 02/03/2023

ABSTRACT

Mung bean is a short-duration ancient legume crop that can be grown as a sole or intercrop for grain and green manure in all three crop seasons, including winter, summer and rainy in various regions of the country. MYMV (Mungbean Yellow Mosaic Virus) disease is one of the most destructive diseases transmitted by whitefly (*Bemisia tabaci* Genn) persistently. Weather parameters have no direct effect on MYMV, but affect the whitefly population. Field experiments were conducted for three consecutive years from 2019-2021 as to know the effect of weather parameters on the whitefly population. Development of whitefly population and MYMV disease severity were recorded in mungbean crops at 7 days interval starting from 20 days after sowing

*Corresponding author: E-mail: vermapreety5926@gmail.com, vermapreety5926@hau.ac.in;

(DAS). The present study showed that whitefly population builds up starting in July and reached to maximum at end of August to early September. There are various factors influencing the build-up of whitefly population viz., maximum temperature $36\pm 2^{\circ}\text{C}$, minimum temperature $25\pm 2^{\circ}\text{C}$, morning relative humidity more than 90 percent, more sunshine hours and no rainfall prevailing during that period. Whitefly population had positive significant correlation with maximum temperature and bright sunshine hours while evening relative humidity and rainfall were recorded as negatively significant. Three-year data is very helpful in guiding the farmers for timely and preventive sprays based on existing infections and the population of whitefly in a season.

Keywords: Persistent manner; MYMV disease; *Bemisia tabaci*; bright sunshine hours; whitefly; preventive spray.

1. INTRODUCTION

“Mung bean (*Vigna radiata* (L.) Wilczek) is a very ancient annual crop that originated in India or the Indo- Burmese region, which is commonly known as green gram, green bean, moong, mash bean, golden gram and green soy. Mung bean (*Fabaceae*) is well suited for a large number of cropping systems. More than 90% of world’s mungbean production comes from Asia” [1]. “Mungbean is a short-duration legume crop, which can be grown in varying environmental conditions, such as in all three crop seasons viz., kharif, summer and rabi in different parts of the country, as sole or intercrop for grain and green manure” [2]. “The global mungbean area is about 7.3 million ha with global output of about 5.3 million tones. The average grain yields (0.73 t/ha) of mungbean are quite low in India. India is the world’s largest producer of mungbean cultivated over an area of 3.44 million ha with a production of 1.4 million tones and productivity of 406.98 kg/ha in India”. (<http://agropedia.iitk.ac.in>).

“The standard yield of mungbean worldwide is very low (730 kg/ha) and the main reasons are various diseases of mungbean. Out of all diseases MYMV disease is one of the most prevalent and destructive diseases” [3]. “The most vulnerable stage is from 30 to 45 days of sowing” [3]. “MYMV a member of the *Geminiviridae* family, belongs to the genus *Begomovirus* that was identified in 1955 and it was observed that it is not transmitted mechanically but transmitted by whitefly (*Bemisia tabaci* Genn) in persistent manner” [4]. “Mungbean plants are infected at the seeding stage, so it can lead to 85-100% loss in yield” [3]. “MYMV disease can be recognized by its typical symptoms which start appearing on the young leaves of the plant at vegetative stage in form of mild yellow spots” [5]. “These gradually spread all over the upper leaves, which turn to yellow

and ultimately, reduce plant size, number of flowers and pods” [6]. “Affected plants generate fewer flowers and pods, which also develop mottling and remain small and contain fewer, smaller and shrunken seeds” [7].

“Plant disease emergences have dramatically increased recently as a result of global changes, especially concerning trade, host genetic uniformity, and climate change” (Willcoquet *et al.*, 2020). “The Mungbean Yellow Mosaic Virus (MYMV) is prevalent in the major mungbean-growing areas in India. A severe outbreak of MYMV in the southern and northern states of India is currently causing serious concern to mungbean growers and to the industrialist in these regions. In past, different varieties had responded differently to. MYMV disease in the field under different temperatures, relative humidity, sunshine hours, wind speed and rainfall. Many weather-driven epidemiological models have been developed and used to predict plant disease epidemics under variable climates” [8]. The present investigation was conducted to form an epidemiological model which may direct the farmers toward early and preventive spraying based on the level of disease incidence and whitefly population during the season.

2. MATERIALS AND METHODS

The present investigation was conducted in the Research Area, Department of Plant Pathology, CCS Haryana Agricultural University, Hisar during Kharif 2019-2022. CCS Haryana Agricultural University is situated in the semi-arid climate at $29^{\circ} 17' \text{ N}$ latitude and $75^{\circ} 47' \text{ E}$ longitude at an altitude of 215.2 meters above mean sea level in the subtropical climatic zone of India. The mean weekly values of all the weather parameters during the crop-growing season were recorded at the Meteorological Observatory of the Research Farm, CCS Haryana Agricultural University, Haryana, India.

To understand the whitefly population and incidence of MYMV disease, a field experiment was conducted to know the effects of various weather factors on the whitefly population and disease severity for three consecutive years. The correlation analysis was performed among disease severity, whitefly population and weather parameters. Mungbean cultivars MH 421 as resistant, TMB 37 as moderately resistant and SML 1082 as susceptible was grown in a randomized block design (RBD) during kharif seasons (2019-2021) in plots. In India, Kharif season lies from May to September.

2.1 Disease Severity

The disease severity was recorded at 7 days interval starting from 20 DAS (Days after sowing) by using the 0-6 scale in Table 1 [9]. Percent disease index (PDI) was calculated by using the method of McKinney, 1923. The correlation was analysed between PDI with the different weather parameters.

Table 1. Disease severity scale (Ali et al., 2005)

Rating	Category	Severity range
0	Immune	0%
1	Highly resistant	1-10%
2	Moderately resistant	11-25%
3	Tolerant	26-50%
4	Moderately susceptible	51-60%
5	Susceptible	61-70%
6	Highly susceptible	71-100%

The percent disease index (PDI) was calculated by using McKinney's [10] formula:

$$\text{Per cent disease index} = (\text{Sum of all numerical ratings} / \text{No. of plants examined} \times \text{maximum disease rating}) \times 100$$

2.2 Management through Biorationals and Chemicals Approaches

2.2.1 Treatment details

T₁: Malathion 50 EC @ 450ml/ acre

T₂: Dimethoate EC @ 250ml/ acre

T₃: Neem oil @ 5ml/kg as seed treatment + spray @5ml/L

T₄: Salicylic acid @ 150 mg/litre as seed treatment + spray

T₅: Sarpagandha leaves extract @10% as seed priming + spray @100ml/L

T₇: Faba bean seed extract @ 10 % as seed priming + spray @100ml/L

3. RESULTS

3.1 Whitefly Population at Weekly Interval

Whitefly population build up on all three cultivars was higher during August as compared to the rest of months. It was recorded that in early stages of crop growth whitefly population was less and build-up started with advances in crop growth. Whitefly population was observed first during the month of July (initial crop growth stages). Thereafter, the whitefly population increased, except during rainfall and high wind speed periods. Whitefly population was ranged between 1.5-8 adults/leaf/plant with an average of 4 adults/leaf/plant on mungbean (Table 2-4). The whitefly population increased up to 45 DAS that might be due to increase in maximum temperature, sunshine hours and decrease in wind speed. The whitefly population builds up started in August. Maximum average whitefly population was recorded (8.42) on SML 1082 followed by TMB 37 (7.19) and minimum (5.08) was recorded on MH 421 at 36.4°C, no rainfall (0.00 mm), minimum relative humidity (53%) and maximum bright sunshine (7.9h) during *kharif* 2019. Among the three cultivars maximum average population was recorded on SML 1082 during the course of disease development in all the three season.

Whitefly population on all three cultivars exhibited positive significant correlation with bright sunshine hours. SML 1082 exhibited positive significant correlation with bright sunshine hours (0.487) while maximum temperature (0.470) and morning relative humidity (0.417) was recorded as negatively non-significant factor (Table 5). However, evening relative humidity (0.846), wind speed (0.471) and rainfall (0.650) was negatively (non-significant) correlated, while minimum temperature (0.470) was recorded positively non-significant with whitefly population. Almost similar correlation matrix was found in case of TMB 37. MH 421 grown as resistant cultivar against MYMV disease also exhibited the same kind of correlation with weather parameters and whitefly population.

3.2 Disease Severity at Weekly Intervals

The MYMV disease occurred throughout the growing season of mungbean crop. The disease

initiated in the field at 20 days after sowing in the form of vein clearing of leaves on 2 to 3 young leaves followed by yellowing of leaves. The higher terminal PDI was observed on SML 1082 (73.32%) cultivar as compared to TMB 37 (41.66%) and MH 421 (10.55) (Table 6-8). At the time of initiation of disease, the maximum temperature was 38±2°C with more than 85 percent morning relative humidity. As the crop growth, the PDI increased significantly. Slow progress in disease was observed in the weeks which had lower whitefly population due to unfavourable weather conditions, i.e. high wind speed (7.21 km/h), less bright sunshine hours (3.82 h) and less total rainfall (83.7mm). They thrive best under hot and humid conditions and the population also towers with higher temperatures.

The progress in disease was also noticed during post rainfall periods. The maximum PDI was observed in last week of August and first week of September. The highest PDI (73.32) was recorded on SML 1082 at 55 DAS during first week of September.

PDI on three cultivars SML 1082 exhibited positive non-significant correlation with minimum temperature, morning relative humidity and bright sunshine hours. However, maximum temperature, wind speed and rainfall were negatively (non-significant) correlated (Table 9). The evening relative humidity is the only factor that has significant effect on PDI that is highly significant (negative) for all the three cultivars.

Table 2. Effect of weather parameters on whitefly population on three cultivars of mungbean during *kharif* 2019

Time of observation	Whitefly population/ leaf			Temperature (°C)		Relative humidity (%)		Average wind velocity (km/h)	Bright sunshine hours (h)	Total Rainfall (mm)
	MH 421	TMB 37	SML 1082	Max.	Min.	M	E	7.21	3.81	23.3
25-07-2020	1.443	2.083	2.593	38.6	24.2	89	74	5.02	4.4	83.7
1-08-2020	1.140	1.570	2.083	35.4	20	89	72	7.5	7.1	1.1
8-08-2020	2.217	3.891	4.590	36.6	26	83	63	4.97	3.2	14.4
15-08-2020	3.797	6.007	6.897	34.4	23.2	91	68	5.18	7.9	0.0
22-08-2020	5.087	7.197	8.420	36.4	25	83	54	4.7	6.2	27.7
29-08-2020	4.830	5.817	6.900	33.5	25.5	85	57			
CD (P=0.05)	1.117	1.029	0.704							
SE±	0.35	0.32	0.22							

Table 3. Effect of weather parameters on whitefly population on three cultivars of mungbean during *kharif* 2020

Time of observation	Whitefly population/ leaf			Temperature (°C)		Relative Humidity (%)		Average wind velocity (km/h)	Bright sunshine hours (h)	Total Rainfall (mm)
	MH 421	TMB 37	SML 1082	Max.	Min.	M	E	6.7	8.0	38.4
16-07-2020	0.00	0.00	0.00	36.0	26.5	89	64	8.7	5.7	111.1
23-07-2020	1.21	3.17	3.32	34.6	25.9	92	75	4.8	7.3	14.7
30-07-2020	3.43	3.48	4.53	34.5	26.8	87	60	5.9	6.6	0.0
06-08-2020	3.76	3.46	4.82	36.4	27.0	84	66	6.9	6.6	32.8
13-08-2020	4.84	4.53	5.67	35.3	27.5	88	62	6.5	6.7	17.5
20-08-2020	5.13	5.34	6.12	35.3	26.9	90	77	7.7	6.4	11.7
27-08-2020	5.28	5.67	6.28	32.6	25.4	91	76	7.5	6.8	0.0
3-09-2020	5.89	5.91	7.50	33.9	25.6	90	64			

Table 4. Effect of weather parameters on whitefly population on three cultivars of mungbean during *kharif* 2021

Time of observation	Whitefly population/ leaf			Temperature (°C)		Relative humidity (%)		Average wind velocity (km/h)	Bright sunshine hours (h)	Total rainfall (mm)
	MH 421	TMB 37	SML 1082	Max.	Min.	M	E	6.7	8.0	38.4
22-07-2020	0.00	0.00	0.00	26.9	35.9	63	86	4.8	8.2	9.0
29-07-2020	1.24	3.34	3.76	27.4	33.7	76	92	8.4	5.9	56.0
5-08-2020	3.78	3.87	4.53	26.5	31.9	79	94	7.6	6.9	149.9
12-08-2020	4.21	4.21	5.22	26.6	35.6	58	88	4.7	7.2	0.0
19-08-2020	4.99	4.78	5.78	26.3	37.1	54	80	4.9	8.4	0.0
26-08-2020	5.65	5.78	6.58	26.7	34.7	69	86	5.9	3.6	0.5
2-09-2022	5.67	6.21	7.32	25.9	35.9	65	88	7.8	5.8	15.7
9-09-2022	5.93	6.43	5.48	25.8	32.7	75	95	6.9	7.4	100.2

Table 5. Correlation matrix of weather parameters and whitefly population on different varieties of mungbean during *kharif* 2019

S. No	Variety	Temperature (°C)		Relative humidity (%)		Wind speed (km/h)	Sunshine hours (h)	Rainfall (mm)
		Maximum	Minimum	Morning	Evening			
1	MH421	-0.477 ^{NS}	0.507 ^{NS}	-0.450 ^{NS}	-0.884 [*]	-0.549 ^{NS}	0.525 ^S	-0.548 ^{NS}
2	TMB37	-0.469 ^{NS}	0.535 ^{NS}	-0.393 ^{NS}	-0.827 [*]	-0.462 ^{NS}	0.462 ^S	-0.661 ^{NS}
3	SML1082	-0.470 ^{NS}	0.539 ^{NS}	-0.417 ^{NS}	-0.846 [*]	-0.471 ^{NS}	0.487 ^S	-0.650 ^{NS}

NS = Non Significant, * = Significant

Table 6. Effect of weather parameters on percent disease index (PDI) on three cultivars of mungbean during *kharif* 2019

Time of observation	Percentage disease index (PDI)			Temperature (°C)		Relative humidity (%)		Average wind velocity (km/h)	Bright sunshine hours (h)	Total Rainfall (mm)
	MH 421	TMB 37	SML 1082	Max.	Min.	M	E	7.21	3.81	23.3
25-07-2020	1.66	6.10	20.02	38.6	24.2	89	74	5.02	4.4	83.7
1-08-2020	1.98	8.88	33.33	35.4	20	89	72	7.5	7.1	1.1
8-08-2020	4.21	17.73	42.77	36.6	26	83	63	4.97	3.2	14.4
15-08-2020	6.11	26.10	49.44	34.4	23.2	91	68	5.18	7.9	0.0
22-08-2020	8.33	32.77	57.77	36.4	25	83	54	4.7	6.2	27.7
29-08-2020	10.55	41.66	73.32	33.5	25.5	85	57			
CD (P=0.05)	3.95	5.57	8.93							
SE±	(1.23)	(1.74)	(2.80)							

• M: Morning relative humidity, E: Evening relative humidity

Positive relationship was found between whitefly population and disease severity. At initial stages of crop growth, whitefly population and disease incidence were found to be lower but later due to favourable weather conditions increase in whitefly population led to increase in MYMV disease. The incidence started at 20 DAS with varying whitefly population on all three cultivars.

The correlation between whitefly population and MYMV PDI was found positively significant. Table 10 shows that MH 421 (0.958) was positively highly significant and. SML 1082 (0.831) and TMB 37 (0.888) were found positively significant which clearly indicated that vector population directly affected the MYMV disease on mungbean crop.

Table 7. Effect of weather parameters on Percent Disease Index (PDI) on three cultivars of mungbean during kharif 2020

Time of observation	Percentage disease index (PDI)			Temperature (°C)		Relative humidity (%)		Average wind velocity (km/h)	Bright sunshine hours (h)	Total rainfall (mm)
	MH 421	TMB 37	SML 1082	Max.	Min.	M	E	6.7	8.0	38.4
16-07-2020	0	0	0	36.0	26.5	89	64	8.7	5.7	111.1
23-07-2020	0	0	2.14	34.6	25.9	92	75	4.8	7.3	14.7
30-07-2020	5.83	4.17	6.7	34.5	26.8	87	60	5.9	6.6	0.0
06-08-2020	6.67	9.17	13.3	36.4	27.0	84	66	6.9	6.6	32.8
13-08-2020	9.17	16.67	26.7	35.3	27.5	88	62	6.5	6.7	17.5
20-08-2020	13.33	22.50	50.0	35.3	26.9	90	77	7.7	6.4	11.7
27-08-2020	15.00	26.67	70.8	32.6	25.4	91	76	7.5	6.8	0.0
3-09-2020	18.33	29.17	75.8	33.9	25.6	90	64			

• M: Morning relative humidity, E: Evening relative humidity

Table 8. Effect of weather parameters on Percent Disease Index (PDI) on three cultivars of mungbean during kharif 2021

Time of observation	Percentage disease index (PDI)			Temperature (°C)		Relative humidity (%)		Average wind velocity (km/h)	Bright sunshine hours (h)	Total rainfall (mm)
	MH 421	TMB 37	SML 1082	Max.	Min.	M	E	6.7	8.0	38.4
22-07-2020	0	0	2.88	26.9	35.9	63	86	4.8	8.2	9.0
29-07-2020	0	4.15	5.77	27.4	33.7	76	92	8.4	5.9	56.0
5-08-2020	4.72	9.57	7.88	26.5	31.9	79	94	7.6	6.9	149.9
12-08-2020	5.87	12.45	15.66	26.6	35.6	58	88	4.7	7.2	0.0
19-08-2020	10.12	19.64	36.99	26.3	37.1	54	80	4.9	8.4	0.0
26-08-2020	14.77	24.58	48.97	26.7	34.7	69	86	5.9	3.6	0.5
2-09-2022	17.88	28.69	67.74	25.9	35.9	65	88	7.8	5.8	15.7
9-09-2022	19.66	32.19	74.89	25.8	32.7	75	95	6.9	7.4	100.2

• M: Morning relative humidity, E: Evening relative humidity

Table 9. Correlation of weather parameters and PDI on different varieties of mungbean

S.No	Variety	Temperature (°C)		Relative humidity (%)		Wind speed	Sunshine hours	Rainfall
		Maximum	Minimum	Morning	Evening			
1	MH421	0.763*	0.533 ^{NS}	0.469 ^{NS}	-0.882 [*]	-0.575 ^{NS}	0.510*	-0.426 ^{NS}
2	TMB37	0.797*	0.612 ^{NS}	0.446 ^{NS}	-0.872 [*]	-0.591 ^{NS}	0.490*	-0.421 ^{NS}
3	SML1082	0.878*	0.436 ^{NS}	0.473 ^{NS}	-0.865 [*]	-0.622 ^{NS}	0.619*	-0.306 ^{NS}

NS = Non significant, * = Significant

Table 10. Correlation matrix of whitefly population and PDI on three mungbean varieties

S. No.	Variety	Disease severity
1	MH 421	0.958 ^{**}
2	TMB 37	0.888 [*]
3	SML 1082	0.831 [*]

** = Highly significant, * = Significant

3.3 Management of MYMV Disease in Mungbean

The results revealed that all the treatments had significant effect on increased the yield as well as test weight. Test weight of all treatments was significantly higher than the control. In all treated plots significantly lower PDI was recorded during all the periods of observations. Amongst the

chemicals, Dimethoate EC recorded significantly less terminal PDI with maximum increase in yield (58.68) followed by Malathion 50 EC @ 450ml/acre (Table 11). Faba bean seed extract as seed treatment + spray @5ml/L shows maximum percent increase in yield (54.68) followed by the use of Neem oil as seed treatment + spray @5ml/L which is at par with the use of salicylic acid as seed treatment + spray @5ml/L.

In order to guide farmers toward early and preventive spraying based on the degree of disease incidence and whitefly population throughout the season, a biorational approaches in mungbean is very helpful.

4. DISCUSSION

For the sake of epidemiological studies temperature, relative humidity, rainfall, wind velocity and sunshine hours were considered as most important factors. The whitefly population and percent disease index (PDI) was observed throughout the growing season on three mungbean cultivars. The disease initiated at 20 DAS and maximum progress in PDI was noticed on. SML 1082 followed by TMB 37 and MH 421. The highest terminal PDI (75.8%) and whitefly population (7.50) was observed on SML 1082. The maximum whitefly population and PDI were observed during last week of August and early September. This might be due to more favourable weather conditions, i.e. maximum temperature 36±2 °C and low evening relative humidity (<54%) with more sunshine hours during this period. No rainfall prevailing during that period. In previous studies, Khan et al., [11] observed maximum whitefly population at higher temperature (38°C) and relative humidity (66%), which were congenial for prevalence of the

mungbean yellow mosaic disease. Similarly, Singh & Gurah [12] found that hot weather with little or no rainfall was conducive for multiplication of MYMV and *Bemisia tabaci*. Meti et al., [13] observed that decrease in whitefly population was mainly due to higher rainfall and lower minimum temperature, whereas increase in whitefly population was due to higher maximum temperature and no rainfall. Cooler weather and high relative humidity are detrimental to whitefly population [14] and present study is in agreement with his findings.

Both the whitefly population and PDI were found negatively significant with evening relative humidity, whereas all other factors were reported having no significant effect. Earlier, Khan et al., (2012) and Ansar et al., [15] have reported negative correlation between maximum temperature and whitefly population. There was positive significant correlation between whitefly population and sunshine hours, which was also observed by Meti et al., [13]. The high wind speed and more total rainfall have detrimental effect on whitefly population. In present findings, the lesser number of whiteflies were observed in the weeks, which have high wind speed and more rainfall, and a negative correlation was found with wind speed and rainfall. Similarly, Marabi et al., [16] has also recorded significant negative correlation with wind speed and rainfall. Present findings were corroborative with the findings of Ali et al., [17] and Suman et al., [18] who reported that the maximum temperature has a significant negative correlation with. MYMV disease incidence and Khan et al., [11] who established negative correlation of the disease with rainfall and wind speed but there was positive correlation of MYMV disease with maximum temperature.

Table 11. Management of MYMV disease in mungbean

Treatment No.	Treatments	Percent increase in yield	Percent increase in test weight	B:C ratio
T1	Malathion 50 EC @ 450ml/ acre	58.68	17.98	1.96
T2	Dimethoate EC @ 250ml/ acre	52.77	16.32	1.84
T3	Neem oil @ 5ml/kg as seed treatment + spray @5ml/L	46.76	13.56	1.69
T4	Salicylic acid @ 150 mg/litre as seed treatment + spray	44.69	12.87	1.56
T5	Sarpagandha leaves extract as seed treatment + spray @5ml/L	38.76	12.37	1.52
T6	Faba bean seed extract as seed treatment + spray @5ml/L	54.68	16.78	1.86
T7	Control	--	--	1.26

Management of MYMV disease by controlling its vector *Bemisia tabaci*, through use of resistance inducers and biorationals was tried. The present result revealed that suppression of MYMV disease could be achieved through use of biorational approaches and nonconventional chemicals. Dubey et al., [19] observed that application of neem seed kernel extract and foliar spray of neem oil is effective. Reang et al., [20] also used different botanicals and out of them, neem @ 0.2% showed lowest disease incidence (10.49%) and severity (9.58%) in case of virus disease.

Studies conducted in tropical agricultural areas of Latin America such as Venezuela [21-23] and Panama have shown that MYMV disease is also a major constraint to mungbean production in these regions [24-27]. The disease is transmitted by the same vector, *Bemisia tabaci*, and causes similar symptoms in infected plants as in India. In Venezuela, MYMV disease has been reported to cause yield losses ranging from 20% to 80%, depending on the severity of the infection [28]. The disease is more prevalent during the rainy season (May to October) [29-31], and the whitefly population is high [32-33]. Management of MYMV disease in Venezuela involves the use of similar strategies as in India, such as cultural practices, chemical control, and host resistance [34,35].

In Panama, MYMV disease is also a major constraint to mungbean production, especially in the central and western regions of the country (the disease is more prevalent during the dry season (December to April) [36], and the whitefly population is low. Management of MYMV disease in Panama involves the use of similar strategies as in India and Venezuela, but the focus is on host resistance. MYMV disease is a major viral disease that affects mungbean production in India and other tropical regions around the world, including Latin America. Management of MYMV disease in these regions involves the use of several strategies such as cultural practices, chemical control, and host resistance. Host resistance is considered the most effective and sustainable approach for managing MYMV disease in mungbean.

5. CONCLUSION

The whitefly population builds up started in July and reached to maximum at end of August to early September. There are various reasons for build-up of whitefly population viz., maximum

temperature $36\pm 2^{\circ}\text{C}$, minimum temperature $25\pm 2^{\circ}\text{C}$, morning relative humidity more than 90 per cent, more sunshine hours and no rainfall prevailing during that period. For the management of MYMV disease biorationals were used and amongst all Faba bean seed extract @ 150 mg/liter as seed priming and then foliar spray @ 150 mg/L recorded significantly less terminal PDI with maximum percent disease control (54.68%).

ACKNOWLEDGEMENTS

All sort of assistance rendered by the Department of Plant Pathology, CCS Haryana Agricultural University, Hisar and Dr. Vinod Kumar Malik for the above study is gratefully acknowledged without whom this work wouldn't have reached its completion.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Karthikeyan A, Shobhana VG, Sudha M, Raveendran M, Senthil N, Pandiyan M, Nagarajan P. Mungbean Yellow Mosaic Virus (MYMV): A threat to green gram (*Vigna radiata*) production in Asia. International Journal of Pest Management. 2015;60(4):314-324.
Available: <https://doi.org/10.1080/09670874.2014.982230>
2. Dharajiya T, Ravindrababu Y, Pagi NK. Screening of mungbean [*Vigna radiata* (L.) Wilczek] genotypes for resistance against Mungbean Yellow Mosaic Virus (MYMV) under field condition. International Journal of Current Microbiology and Applied Sciences. 2018;7(5):3472-3483.
Available: <https://doi.org/10.20546/ijcmas.2018.705.401>
3. Sudha M, Karthikeyan A, Anusuya P, Ganesh NM, Senthil N, Pandiyan M, Raveendran M, Nagarajan P, Angappan K. Inheritance of resistance to Mungbean Yellow Mosaic Virus (MYMV) in inter and intra specific crosses of mungbean (*Vigna radiata*). American Journal of Plant Sciences. 2013;4:1924-1927.
Available: <http://dx.doi.org/10.4236/ajps.2013.410236>

4. Iqbal U, Iqbal SM, Afzal R, Jamal A, Farooq MA, Zahid A. Screening of mungbean germplasm against Mungbean Yellow Mosaic Virus (MYMV) under field conditions. *Pakistan Journal of Phytopathology*. 2011;23(1):48-51.
5. John NB, Packiaraj D, Pandiyan M, Senthil N. Screening of mungbean (*Vigna radiata* L.) genotypes for resistance to yellow mosaic virus. *Trends in Biosciences*. 2015;8(8):2042-2046.
6. Brown SB, Houghton JD, Hendry GAF. Chlorophyll breakdown. In: *Chlorophylls*, H. Scheer (Ed). CRC Press: Boca Raton, Florida. 1991:465–489.
7. Patel P, Modha K, Vadodariya G, Patel R. Screening of mungbean germplasm for resistance to mungbean yellow mosaic virus under natural condition. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(5):864-867.
8. Datta P, Dasgupta B, Majumder D, Das S. Disease prediction model of foot rot and leaf rot of betelvine (*Piper betle* L.) caused by *Phytophthora parasitica* in West Bengal. SAARC. *Journal of Agriculture*. 2010;8(2): 87-96.
9. Ali S, Khan MA, Habib A, Rasheed S, Iftikhar Y. Management of yellow vein mosaic disease of okra through pesticide/bio-pesticide and suitable cultivars. *International Journal of Agriculture & Biology*. 2005;7(1):145-147.
10. Mckinney HH. Influence of soil temperature and moisture on infection of wheat seedling by *Helminthosporium sativum*. *Journal of Agricultural Research*. 1923;26: 195-217.
11. Khan MA, Haider S, Ali Yashir. Evaluation of mungbean germplasm for resistance against mungbean yellow mosaic virus and whitefly population in relation to epidemiological factors. *Agricultural Research and Technology*. 2018;18(3): 001-008.
DOI: 10.19080/ARTOAJ.2018.18.556058
12. Singh RA, Gurha SN. Influence of cropping seasons on the incidence of yellow mosaic disease in mungbean genotypes. *Indian Journal of Pulses Research*. 1994;7(12): 206-208.
13. Meti M, Kenganal M. Epidemiology of Mungbean Yellow Mosaic Virus (MYMV) in relation to whitefly dynamics and weather parameters. *International journal of Current Microbiology and Applied Sciences*. 2018; 7(9):2368-2375.
Available: <https://doi.org/10.20546/ijcmas.2018.709.294>
14. Sharma S, Yogesh K, Shyam S. Influence of abiotic weather parameters on population dynamics of whitefly, *Bemisia tabaci* (Genn) on cotton. *Journal of Cotton Research and Development*. 2014;28(2) :286-288.
15. Ansar M, Saha T, Sarkhel S, Bhagat AP. Epidemiology of okra yellow vein mosaic disease and its interaction with insecticide modules. *Trends in Biosciences*. 2014;7 (24):4157-4160.
16. Marabi RS, Das SB, Bhowmick AK, Pachori R, Vibha H, Sharma HL. Seasonal population dynamics of whitefly in soybean. *Journal of Entomology and Zoology Studies*. 2017;5(2):169-173.
17. Ali S, Khan MA, Zeshan MA, Habib A, Haider MS. Characterization of conducive environmental conditions for mungbean yellow mosaic virus disease incidence on mungbean germplasm. *Pakistan Journal of Phytopathology*. 2015;27(1): 27-30.
18. Suman S, Kumar M, Sharma VK, Kumar H. Correlation of weather parameters with incidence of Mungbean Yellow Mosaic Virus (MYMV) disease in mungbean (*Vigna radiata* (L. Wilczek). *International Journal of Agricultural Statistical Sciences*. 2016;12:109-115.
19. Dubey NK, Shukla R, Kumar A, Singh P, Prakash B. Global scenario on the application of natural products in integrated pest management programmes. In: Dubey NK, editor. *Natural products in plant pest management*, Wallingford: CAB International. 2011;1:1-20.
20. Reang D, Devi MR, Dhar A, Nath PS. Comparative evaluation of botanicals and chemical insecticides and their suitable use for the management of Yellow Mosaic Virus (YMV) of mungbean. *International Journal of Current Microbiology and Applied Sciences*. 2018;7(9):2319-7706.
Available: <https://doi.org/10.20546/ijcmas.2018.709.255>.
21. Olivares B, Hernández R, Arias A, Molina JC, Pereira Y. Identificación de zonas agroclimáticas potenciales para

- producción de cebolla (*Allium cepa* L.) en Carabobo, Venezuela. *Journal of the Selva Andina Biosphere*. 2018;6(2):70-82.
Available:http://www.scielo.org.bo/pdf/jsab/v6n2/v6n2_a03.pdf
22. Hernández R, Coelho R, Molina JC, Pereira Y, Orlando B. Analysis of climate types: Main strategies for sustainable decisions in agricultural areas of Carabobo, Venezuela. *Scientia Agropecuaria*. 2018;9(3):359–369.
DOI: 10.17268/sci.agropecu.2018.03.07
 23. Orlando O, Hernández R, Arias A, Molina JC, Pereira Y. Zonificación agroclimática del cultivo de maíz para la sostenibilidad de la producción agrícola en Carabobo, Venezuela. *Revista Universitaria de Geografía*. 2018;27(2):139-159.
Available:<https://n9.cl/l2m83>
 24. Montenegro E, Pitti J, Olivares B. Adaptation to climate change in indigenous food systems of the Teribe in Panama: training based on CRISTAL 2.0. *Luna Azul*. 2021a;51(2):182-197.
Available:<https://n9.cl/qwvwz>
 25. Montenegro E, Pitti J, Olivares B. Identificación de los principales cultivos de subsistencia del Teribe: Un estudio de caso basado en técnicas multivariadas. *Idesia*. 2021b;39(3):83-94.
Available:<http://dx.doi.org/10.4067/S0718-34292021000300083>
 26. Bertorelli M, Olivares BO. Population fluctuation of *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) in sorghum cultivation in Southern Anzoategui, Venezuela. *Journal of Agriculture University of Puerto Rico*. 2020;104(1):1-16.
Available:<https://doi.org/10.46429/jaupr.v104i1.18283>
 27. Olivares B, Hernández R. Ecoterritorial sectorization for the sustainable agricultural production of potato (*Solanum tuberosum* L.) in Carabobo, Venezuela. *Agricultural Science and Technology*. 2019;20(2):339-354.
Available:https://doi.org/10.21930/rcta.vol20_num2_art:1462
 28. Demey JR, Olivares B, Zingaretti ML, Demey Zambrano JA. Tipificación de los sistemas de producción agrícola y la percepción de la variabilidad climática en Anzoátegui, Venezuela. *Revista FAVE - Ciencias Agrarias*. 2016;15(2):39-50.
Available:<https://doi.org/10.14409/fa.v15i2.6587>
 29. Rey JC, Lobo D, Gómez JA, Olivares B, Landa B.. Impacto del cambio climático en zonas bananeras de la Región Central de Venezuela: El futuro de los bananos en un escenario hídrico incierto. En: Chica Pérez, A. F. y Mérida García, J. (Eds). *Creando Redes Doctorales: Investiga y Comunica*. Córdoba, España: UCO Press. Editorial Universidad de Córdoba. 2019: 367-370.
 30. Parra R, Cortez A, Rodríguez MF, Orlando B. Patrones de homogeneidad pluviométrica en estaciones climáticas del estado Anzoátegui, Venezuela. *Revista Multiciencias*. 2012;12(Extraordinario):11-17.
Available:<https://n9.cl/xbslq>
 31. Cortez A, Olivares B, Parra R, Lobo D, Rodríguez MF, Rey JC. Descripción de los eventos de sequía meteorológica en localidades de la cordillera central, Venezuela. *Ciencia, Ingenierías y Aplicaciones*. 2018;1(1):22-44.
Available:<http://dx.doi.org/10.22206/cyap.2018.viil.pp23-45>
 32. Hernández R, Pereira Y, Molina JC, Coelho R, Olivares BY Rodríguez K. Calendario de siembra para las zonas agrícolas del estado Carabobo en la República Bolivariana de Venezuela. Sevilla, Spain, Editorial Universidad Internacional de Andalucía. 2017:247.
 33. Hernandez R, Arias A, Molina JC, Pereira Y, Campos O. Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. *Idesia*. 2020;38(2):95-102.
Available:<http://dx.doi.org/10.4067/S0718-34292020000200095>
 34. Guevara E, Demey J, Campos O. Utilización de bioindicadores climáticos en sistemas de producción agrícola del estado Anzoátegui, Venezuela. *Revista Multiciencias*. 2012;12(2):136-145.
Available:<https://n9.cl/ak22r>
 35. Olivares B. La sequía meteorológica en territorios agrícolas de Venezuela: Un análisis temporal del fenómeno meteorológico y su impacto en la agricultura venezolana. Saarbrücken,

- Germany, Academic Spanish Editorial. 2017;145.
36. Pitti J, Olivares B, Montenegro E. The role of agriculture in the Changuinola district: A case of applied economics in Panama. Tropical and Subtropical Agroecosystems. 2021;25(1):1-11.
Available:<https://www.revista.ccba.uady.mx/ojs/index.php/TSA/article/view/3815>

© 2023 Verma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/97085>